

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1986

[Full Table of Content](#)



## 86/R/CA/1 Anti-feedants - Cabbages

### Rothamsted Research

Rothamsted Research (1987) *86/R/CA/1 Anti-feedants - Cabbages* ; Yields Of The Field Experiments 1986, pp 296 - 297 - DOI: <https://doi.org/10.23637/ERADOC-1-36>

86/R/CA/1

CABBAGES

ANTI-FEEDANTS

Object: To study the effects of anti-feedants on insect pests and on yields of cabbages - Long Hoos III 5.

Sponsor: D.C. Griffiths.

Design: 4 randomised blocks of 8 plots.

Whole plot dimensions: 2.4 x 5.0.

Treatments:

CHEMICAL	Chemical foliar sprays:
NONE	None (triplicated)
AJUGA EX	Ajuga extract at 0.20 kg
PIPERINE	Piperine at 0.20 kg
POLYGODI	Polygodial at 0.10 kg
E B FARN	Undecyl adduct of (E)-beta-farnesene at 1.0 kg
DELTAMET	Deltamethrin at 0.01 kg

NOTE: Chemical treatments were applied ten times with an electrostatic sprayer in 10 l on 27 June, 1986, 3 July, 9 July, 8 Aug, 14 Aug, 20 Aug, 28 Aug, 5 Sept, 11 Sept and 21 Sept.

Basal applications: Manures: Muriate of potash at 520 kg. (10:10:15 + 4.5 Mg) at 2700 kg. Weedkillers: Paraquat at 0.84 kg ion in 220 l. Clopyralid at 0.10 kg in 220 l and on a second occasion at 0.20 kg in 220 l with the desmetryne. Desmetryne at 0.38 kg in 220 l and on a second occasion at 0.36 kg with the clopyralid.

Seed: Brunswick, sown at 8 kg, in rows 48 cm apart. Plants were thinned to 23 cm apart within the rows on 29 July.

Cultivations, etc.: - Muriate of potash applied: 11 Nov, 1985. Deep-tine cultivated: 19 Dec and again on 19 May, 1986. NPK Mg applied: 28 May. Spring-tine cultivated twice, seed sown, paraquat applied: 2 June. Clopyralid applied: 7 July. Desmetryne applied: 16 July. Clopyralid with desmetryne applied: 20 Aug. Harvested by hand: 26 Nov. Previous crops: S. barley 1984, potatoes 1985.

NOTE: Flea beetle and caterpillar damage were assessed on four occasions during the season.

86/R/CA/1

FRESH YIELD TONNES/HECTARE

\*\*\*\*\* TABLES OF MEANS \*\*\*\*\*

CHEMICAL	NONE	AJUGA	EX	PIPERINE	POLYGODI	E B	FARN	DELTAMET	MEAN	
	98.4		94.0		94.5		87.1	84.9	104.4	95.0

\*\*\*\*\* STANDARD ERRORS OF DIFFERENCES OF MEANS \*\*\*\*\*

TABLE	CHEMICAL
-----	-----
SED	4.45 MIN REP
	3.63 MAX-MIN

	CHEMICAL
MAX-MIN	NONE V ANY OF THE REMAINDER
MIN REP	ANY OF THE REMAINDER

\*\*\*\*\* STRATUM STANDARD ERRORS AND COEFFICIENTS OF VARIATION \*\*\*\*\*

STRATUM	DF	SE	CV%
BLOCK.WP	23	6.29	6.6
PLOT AREA HARVESTED	0.00072		