

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1987

[Full Table of Content](#)

ROTHAMSTED RESEARCH  
Rothamsted Experimental Station  
Harpenden  
Herts  
SG8 5LR  
ENGLAND  
UK

The Rothamsted Research Library is a registered charity and is a member of the British Library. It is a registered charity and is a member of the British Library. It is a registered charity and is a member of the British Library.

Printed: Northampton, 1988

## 87/R/CS/302 Eyespot Resistance to Mbc - W. Wheat

### Rothamsted Research

Rothamsted Research (1988) *87/R/CS/302 Eyespot Resistance to Mbc - W. Wheat* ; Yields Of The Field Experiments 1987, pp 116 - 116 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/CS/302

EYESPOT RESISTANCE TO MBC

Object: To study the development of resistance to MBC fungicides in eyespot and the ability of resistant strains to survive, spread and infect - Meadow.

Sponsors: G.L. Bateman, B.D.L. Fitt.

The third years, w. wheat.

For previous years see 85-86/R/CS/302.

Design: 2 randomised blocks of 4 plots split into 6.

Whole plot dimensions: 12.0 x 24.0.

Treatments: All combinations of:-

Whole plots

1. FUNGCIDE      Fungicides applied cumulatively to 1985 and 1986 treatments:

NONE	None
CARB	Carbendazim at 0.25 kg
PRO	Prochloraz at 0.40 kg
CARB+PRO	Carbendazim at 0.15 kg + prochloraz at 0.40 kg

Sub plots

2. EYE INOC      Eyespot inoculum, applied in first year only:

NATURAL	Natural background population (duplicated)
W 19R 1S	Inoculated with wheat strains in proportion 19 resistant to one sensitive
W 1R 19S	As above but one resistant to 19 sensitive
R 19R 1S	Inoculated with rye strains, 19 resistant to one sensitive
R 1R 19S	As above but one resistant to 19 sensitive

NOTES: (1) Fungicide treatments were applied in 500 l on 12 Nov, 1986 repeated in 200 l on 14 Apr, 1987.

(2) The eyespot inoculum was colonised on oat seed and this was broadcast in October, 1984.

Basal applications: Manures: 'Nitram' at 590 kg. Weedkillers: Paraquat at 0.60 kg ion in 200 l. Isoproturon at 2.5 kg with clopyralid at 0.07 kg, bromoxynil at 0.34 kg and mecoprop at 2.5 kg in 200 l.

Seed: Avalon, sown at 180 kg.

Cultivations, etc.: - Heavy spring-tine cultivated twice: 27 Aug, 1986. Paraquat applied: 18 Sept. Discd, rotary harrowed, seed sown: 30 Sept. Remaining weedkillers applied: 16 Apr, 1987. N applied: 17 Apr. Combine harvested: 1 Sept.

NOTE: Yields were not taken. Eyespot and sharp eyespot were assessed in April and July. Eyespot was characterized according to type and MBC resistance.