

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

ARC, Institute of Arable Crops Research
Rothamsted Experimental Station
Harpenden
Herts
SG8 5LR
ENGLAND
UK

The copyright in this document is owned by the Rothamsted Research Ltd
and is a part of the Rothamsted Research Archives. It is made available
online as a service to the public. It is not to be used for any other purpose
without the prior written permission of Rothamsted Research.

Printed: 2016-06-06
Rothamsted 2016

87/R/CS/312 Straw Decomposition - W. Wheat

Rothamsted Research

Rothamsted Research (1988) *87/R/CS/312 Straw Decomposition - W. Wheat* ; Yields Of The Field Experiments 1987, pp 126 - 127 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/CS/312

STRAW DECOMPOSITION

Object: To test the effects of two basidiomycetes on the decomposition of wheat straw from a preceding crop and on the establishment and yield of a following crop - West Barnfield I.

Sponsor: S.H.T. Harper.

The third year, w. wheat.

For previous years see 85-86/R/CS/312.

Design: 4 randomised blocks of 4 plots.

Whole plot area: 4.5 x 12.0.

Treatments: All combinations of treatments applied to chopped straw in the field, cumulative to applications in the first year:

- | | |
|----------------|--|
| 1. TREATMNT[1] | Treatment one: |
| NONE | None |
| BASID 1 | Basidiomycete 1, cumulative to whey at 15 kg in 1985 and 1986 |
| 2. TREATMNT[2] | Treatment two: |
| NONE | None |
| BASID 2 | Basidiomycete 2, cumulative to a fungal accelerator in 1985 and 1986 |

- NOTES: (1) Basidiomycetes 1 and 2 were naturally occurring fungi found in soil at Rothamsted and Woburn respectively.
(2) The basidiomycete fungus was colonised on wheat seed and this was spread on the surface at 1 seed per square cm on 25 Sept, 1986.
(3) Straw was chopped by a trailed straw chopper and incorporated to a depth of about 10 cm by a rotary grubber.

Basal applications: Manures: 'Nitram' at 130 kg and later at 590 kg.
Weedkillers: Paraquat at 0.60 kg ion in 200 l. Isoproturon at 2.5 kg in 200 l. Clopyralid at 0.05 kg and bromoxynil at 0.24 kg with mecoprop at 1.8 kg in 200 l. Fungicides: Triadimefon at 0.12 kg and carbendazim at 0.25 kg in 200 l. Prochloraz at 0.40 kg and carbendazim at 0.15 kg in 200 l. Propiconazole at 0.12 kg in 200 l. Propiconazole at 0.12 kg with carbendazim at 0.25 kg and maneb at 1.5 kg in 200 l. Insecticide: Pirimicarb at 0.14 kg in 200 l. Molluscicide: Methiocarb at 0.22 kg.

Seed: Mission, sown at 190 kg.

87/R/CS/312

Cultivations, etc.:— Straw chopped: 21 Aug, 1986. Cultivated with rotary grubber: 26 Sept. Methiocarb applied: 29 Sept. Paraquat applied: 3 Oct. Rotary harrowed: 6 Oct. Seed sown: 7 Oct. First N applied: 19 Mar, 1987. Isoproturon applied: 31 Mar. Triadimefon and carbendazim applied: 14 Apr. Second N, clopyralid, bromoxynil and mecoprop applied: 18 Apr. Prochloraz and carbendazim applied: 27 Apr. Propiconazole applied: 28 May. Propiconazole with carbendazim and maneb applied, insecticide applied separately: 23 June. Combine harvested: 19 Aug.

NOTE: Samples of straw were taken throughout the season for observations on the rate of decomposition.

GRAIN TONNES/HECTARE

***** Tables of means *****

TREATMNT[2] TREATMNT[1]	NONE	BASID 2	Mean
NONE	6.85	6.54	6.70
BASID 1	6.68	6.28	6.48
Mean	6.76	6.41	6.59

*** Standard errors of difference's of means ***

Table	TREATMNT[1]	TREATMNT[2]	TREATMNT[1] TREATMNT[2]
s.e.d.	0.258	0.258	0.364

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	9	0.515	7.8

GRAIN MEAN DM% 83.2

PLOT AREA HARVESTED 0.00288