

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1987

[Full Table of Content](#)

ARC, Institute of Arable Crops Research  
Rothamsted Experimental Station  
Harpenden  
Herts  
AL5 2JQ  
UK  
© Rothamsted Research 1988  
All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or by any information storage or retrieval system, without prior permission in writing from Rothamsted Research.  
Printed: Northampton, 1988

## 87/R/CS/320 Comparison of Combinable Crops - W. Wheat

### Rothamsted Research

Rothamsted Research (1988) *87/R/CS/320 Comparison of Combinable Crops - W. Wheat* ; Yields Of The Field Experiments 1987, pp 132 - 134 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/CS/320

COMPARISON OF COMBINABLE CROPS

Object: To compare yields and other attributes of a range of combinable crops and to study their effects on a following crop of w. wheat - Long Hoos VI/VII 5.

Sponsors: J. McEwen, D.P. Yeoman, A.E. Johnston, R.J. Darby.

The second year, w. wheat, s. wheat.

For previous year see 86/R/CS/320.

Design: 3 randomised blocks of 10 plots split into 2.

Whole plot dimensions: 2.5 x 8.0.

Treatments: All combinations of:-

Whole plots

1. PREVCROP	Crops in 1986:
W BEANS	W. field beans, <i>Vicia faba</i>
W OATS	W. oats
W RAPE	W. oilseed rape
W PEAS	W. peas, <i>Pisum sativum</i>
W WHEAT	W. wheat
S BEANS	S. field beans, <i>Vicia faba</i>
S LUPINS	S. lupins, <i>Lupinus albus</i>
S PEAS	S. peas, <i>Pisum sativum</i>
SNFLOWER	Sunflower
FALLOW	Fallow

Sub plots

2. SPRING N	Nitrogen fertilizer applied on 10 Apr, 1987:
0	None
N	Applied, amount depending on quantity in crop and soil in spring

NOTES: (1) Amounts of N applied (kg N) as 'Nitro-Chalk' were:

After PREVCROP	W RAPE, S PEAS, FALLOW	190
	W BEANS	200
	S BEANS, SUNFLOWERS	210
	W OATS, W WHEAT	230
	W PEAS, S LUPINS	240

(2) W. wheat after PREVCROP S LUPINS failed and was resown to s. wheat.

87/R/CS/320

Standard applications:

After all treatments except after lupins: Weedkillers: Terbutryne at 2.8 kg with paraquat at 0.40 kg ion in 220 l. Isoproturon at 2.1 kg (2.5 kg after s. beans and sunflowers) with mecoprop at 2.8 l (2.0 l after s. beans and sunflowers) in 220 l. Cyanazine at 0.35 kg, clopyralid at 0.06 kg and mecoprop at 1.7 l in 220 l applied with the fungicides. Fungicides: Prochloraz at 0.40 kg, carbendazim at 0.15 kg.

After lupins: Weedkillers: Bentazone at 0.80 kg, dichlorprop at 1.1 kg and MCPA at 0.64 kg in 220 l applied with the fungicide. Fungicide: Tridemorph at 0.52 Kg.

Seed: W. wheat: Avalon, sown at 200 kg.  
S. wheat: Wembley, sown at 180 kg.

Cultivations, etc.:-

After w. beans, w. oats, w. rape, w. peas, w. wheat and s. peas. Deep-tine cultivated, rotary cultivated: 13 Aug, 1986 (after w. oats, w. rape, w. peas and fallow only). Shallow-tine cultivated, rotary cultivated: 15 Aug (after w. wheat only). Shallow-tine cultivated: 1 Sept (after s. peas only). Spring-tine cultivated: 11 Sept (after w. beans only). Rotary cultivated: 1 Sept (11 Sept after w. beans). Power harrowed, seed sown, rolled: 23 Sept. Terbutryne and paraquat applied: 24 Sept. Isoproturon and mecoprop applied: 29 Oct and 2 Apr, 1987. Cyanazine, clopyralid, mecoprop and fungicides applied: 16 Apr. Combine harvested: 1 Sept.

After s. beans and sunflowers: Spring-tine cultivated, rotary cultivated: 2 Oct, 1986 (after s. beans only). Ploughed: 13 Oct. Spring-tine cultivated, seed sown, rolled, terbutryne and paraquat applied: 14 Oct. Isoproturon and mecoprop applied: 2 Apr, 1987. Cyanazine, clopyralid, mecoprop, prochloraz and carbendazim applied: 16 Apr. Combine harvested: 1 Sept.

After lupins: W. wheat sown, spring-tine cultivated: 4 Dec, 1986. Spring-tine cultivated, s. wheat sown, rolled: 31 Mar, 1987. Bentazone, dichlorprop, MCPA and tridemorph applied: 19 May. Combine harvested: 10 Sept.

- NOTES: (1) Take-all was assessed in mid July.  
(2) Amounts of ammonium and nitrate nitrogen in the soil were measured in autumn and late winter.  
(3) N contents of grain were measured.  
(4) The wrong rate of SPRING N was applied to two plots, those with treatment combinations S PEAS 0 and SNFLOWER N. Estimated values were used in the analysis.

87/R/CS/320

W. WHEAT (S. WHEAT AFTER S. LUPINS)

GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

SPRING N PREVCROP	0	N	Mean
W BEANS	3.53	8.10	5.82
W OATS	2.18	8.13	5.16
W RAPE	4.10	8.63	6.37
W PEAS	3.28	7.83	5.56
W WHEAT	2.50	6.73	4.62
S BEANS	3.98	7.89	5.93
S LUPINS	3.26	4.93	4.09
S PEAS	3.43	7.88	5.65
SNFLOWER	2.71	7.78	5.24
FALLOW	5.28	7.94	6.61
Mean	3.42	7.59	5.50

\*\*\* Standard errors of differences of means \*\*\*

Table	PREVCROP	SPRING N	PREVCROP SPRING N
s.e.d.	0.349	0.188	0.546
Except when comparing means with the same level(s) of PREVCROP			0.593

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	18	0.428	7.8
BLOCK.WP.SP	18	0.726	13.2

GRAIN MEAN DM% 83.2

PLOT AREA HARVESTED 0.00055