Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1987



Full Table of Content

87/W/CS/322 Effects of Globodera Pallida - Potatoes

Rothamsted Research

Rothamsted Research (1988) 87/W/CS/322 Effects of Globodera Pallida - Potatoes; Yields Of The Field Experiments 1987, pp 135 - 137 - DOI: https://doi.org/10.23637/ERADOC-1-37

87/W/CS/322

EFFECTS OF GLOBODERA PALLIDA

Object: To study the residual effects of a range of potato varieties differing in susceptibility to Globodera pallida and of nematicides on the numbers of Globodera pallida and on the yield of a susceptible variety grown in the second year - Far Field II.

Sponsor: A.G. Whitehead.

The second year, potatoes.

For first year see 86/W/P/3.

Design: 3 randomised blocks of 32 plots.

Whole plot dimensions: 3.0 x 4.57.

Treatments: All combinations of:-

1. VARIETY[86] Varieties in 1986, all Pentland Crown in 1987:

12290 AF 12290 AF20 A 25/11 A 25/11 Cara CARA CROMWELL Cromwell. Desiree DESIREE DIANA Diana HEATHER Heather KIRSTY Kirsty MARFONA Marfona MORAG Morag Maris Piper PIPER Romano ROMANO Sante SANTE ZB 35/29 ZB 35/29

0.0

plus four extra treatments:

VAR NEM[86] Varieties and nematicides in 1986, Pentland Crown, no nematicide in 1987:

CARA ALD

Cara with slow-release aldicarb at 5.6 kg

CARA CAR

PIPR ALD

Cara with carbofuran at 5.6 kg

Maris Piper with slow-release aldicarb at 5.6 kg

PIPR CAR Maris Piper with carbofuran at 5.6 kg

Basal applications: Manures: (10:10:15+4.5 Mg) at 2290 kg. Weedkiller: Linuron at 1.6 kg in 200 l. Fungicides: Mancozeb at 1.4 kg in 200 l on four occasions with the pirimicarb on the second occasion. Fentin hydroxide at 0.28 kg in 200 l on two occasions. Insecticide: Pirimicarb at 0.14 kg. Desiccant: Diquat at 0.80 kg ion in 200 l.

87/W/CS/322

Cultivations, etc.:- Deep-tine cultivated: 30 Jan, 1987 and 16 Feb. NPK Mg applied: 22 Apr. Rotary cultivated and potatoes planted: 28 Apr. Rotary ridged: 15 May. Weedkiller applied: 18 May. Mancozeb applied: 24 June, 26 July, 5 Aug. Mancozeb and pirimicarb applied: 8 July. Fentin hydroxide applied: 18 Aug, 4 Sept. Desiccant applied: 18 Sept. Haulm mechanically destroyed: 1 Oct. Lifted: 7 Oct.

NOTE: Soil samples were taken before nematicides were applied for cyst and egg counts of Globodera pallida.

TOTAL TUBERS TONNES/HECTARE

***** Tables of means *****

OXAMYL[86] VARIETY[86]	0.0	5.6	Mean
12290 AF	45.7	58.7	52.2
A 25/11	55.1	59.5	57.3
CARA	36.9	55.2	46.0
CROMWELL	46.8	57.7	52.3
DESIREE	42.3	52.3	47.3
DIANA	37.5	56.7	47.1
HEATHER	52.0	62.7	57.3
KIRSTY	42.9	54.7	48.8
MARFONA	45.1	55.0	50.1
MORAG	53.1	58.5	55.8
PIPER	44.3	54.2	49.2
ROMANO	46.1	59.1	52.6
SANTE	46.6	55.1	50.8
ZB 35/29	54.5	55.1	54.8
Mean	46.3	56.8	51.5

VAR NEM[86] CARA ALD CARA CAR PIPR ALD PIPR CAR Mean 55.8 42.6 61.1 43.5 50.8

Grand mean 51.4

*** Standard errors of differences of means ***

Table VAR NEM[86] VARIETY[86] OXAMYL[86] VARIETY[86] OXAMYL[86] s.e.d. 3.80 2.69 1.02 3.80

***** Stratum standard errors and coefficients of variation *****

Stratum d.f. s.e. cv% BLOCK.WP 62 4.66 9.1

87/W/CS/322

PERCENTAGE WARE 4.44CM (1.75 INCH) RIDDLE

***** Tables of means *****

OXAMYL[86]	0.0	5.6	Mean
VARIETY[86]	05.4	05.0	05.0
12290 AF	95.4	95.0	95.2
A 25/11	94.9	95.3	95.1
CARA	93.2	95.7	94.4
CROMWELL	96.0	95.4	95.7
DESIREE	94.6	96.0	95.3
DIANA	92.6	95.9	94.2
HEATHER	96.0	95.8	95.9
KIRSTY	94.2	95.0	94.6
MARFONA	94.5	95.3	94.9
MORAG	96.0	96.2	96.1
PIPER	94.0	95.8	94.9
ROMANO	95.8	96.7	96.2
SANTE	95.2	95.5	95.3
ZB 35/29	96.0	96.5	96.2
Mean	94.9	95.7	95.3

VAR NEM[86] CARA ALD CARA CAR PIPR ALD PIPR CAR 95.5 94.2 96.2 94.5 95.1

Grand mean 95

95.3

PLOT AREA HARVESTED 0.00070