

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1987

[Full Table of Content](#)

ARC, Institute of Arable Crops Research  
Rothamsted Experimental Station  
Harpenden  
Herts  
AL5 2JQ  
UK  
Tel: 0494 473941  
Fax: 0494 473942  
Email: [arc@rothamsted.ac.uk](mailto:arc@rothamsted.ac.uk)  
Web: <http://www.rothamsted.ac.uk>  
© Rothamsted Research 1988

## 87/R/WW/5 Aphicide, N and Fungicide - W. Wheat

### Rothamsted Research

Rothamsted Research (1988) *87/R/WW/5 Aphicide, N and Fungicide - W. Wheat* ; Yields Of The Field Experiments 1987, pp 162 - 163 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/WW/5

WINTER WHEAT

APHICIDE, N AND FUNGICIDE

Object: To determine the economic thresholds for cereal aphids with different levels of inputs - Delafield.

Sponsor: N. Carter.

Design: 3 randomised blocks of 12 plots.

Whole plot dimensions: 3.0 x 12.0.

Treatments: All combinations of:-

1. APHICIDE                Aphicides (applied in 200 l):  

NONE	None
PIRIMICA	Pirimicarb applied at 0.14 kg on 23 June, 1987, 3 July and 16 July
2. N RATE                 Nitrogen fertilizers (kg N) as 'Nitram' on 14 Apr, 1987:  

80	
120	
160	
3. FUNGICIDE             Fungicides:  

NONE	None
31+39+59	Fungicide sprays at growth stage 31, 39, 59: G.S. 31 - Prochloraz at 0.40 kg and carbendazim at 0.15 kg in 380 l on 6 May, 1987 G.S. 39 - Propiconazole at 0.12 kg in 200 l on 28 May G.S. 59 - Propiconazole at 0.12 kg with carbendazim at 0.25 kg and maneb at 1.5 kg in 200 l on 23 June

Basal applications: Weedkillers: Clopyralid at 0.07 kg and bromoxynil at 0.34 kg with mecoprop at 2.5 kg in 200 l. Glyphosate at 1.4 kg in 200 l. Growth regulator: Chlormequat at 1.6 kg in 200 l.

Seed: Avalon, sown at 200 kg.

Cultivations, etc.:- Rotary harrowed, seed sown: 6 Nov, 1986. Clopyralid, bromoxynil and mecoprop applied: 23 Apr, 1987. Growth regulator applied: 6 May. Glyphosate applied: 17 Aug. Combine harvested: 1 Sept. Previous crops: W. wheat 1985, potatoes 1986.

NOTE: Aphids were counted from early June until late July. Plant samples were taken at anthesis for dry weight measurements. Disease assessments were made in late June and late July. Components of yield were measured.

87/R/WW/5

GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

N RATE	80	120	160	Mean
APHICIDE				
NONE	6.62	6.96	7.21	6.93
PIRIMICA	6.73	7.27	7.47	7.16
Mean	6.68	7.12	7.34	7.04

FUNGCIDE	NONE	31+39+59	Mean
APHICIDE			
NONE	6.40	7.46	6.93
PIRIMICA	6.66	7.65	7.16
Mean	6.53	7.56	7.04

FUNGCIDE	NONE	31+39+59	Mean
N RATE			
80	6.39	6.96	6.68
120	6.57	7.66	7.12
160	6.64	8.04	7.34
Mean	6.53	7.56	7.04

	N RATE	80	120	160			
APHICIDE	FUNGCIDE	NONE	31+39+59	NONE	31+39+59	NONE	31+39+59
NONE		6.25	7.00	6.50	7.42	6.46	7.95
PIRIMICA		6.54	6.92	6.65	7.90	6.81	8.13

\*\*\* Standard errors of differences of means \*\*\*

Table	APHICIDE	N RATE	FUNGCIDE	APHICIDE N RATE
s.e.d.	0.124	0.152	0.124	0.215

Table	APHICIDE FUNGCIDE	N RATE FUNGCIDE	APHICIDE N RATE FUNGCIDE
s.e.d.	0.176	0.215	0.304

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	22	0.373	5.3

GRAIN MEAN DM% 83.9

PLOT AREA HARVESTED 0.00258