

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

ARC, Institute of Arable Crops Research
Rothamsted Experimental Station
Harpenden
Herts
AL5 2JQ
UK
The copyright in this document is owned by the Rothamsted Research Ltd.
This document is published on the condition that its use for any other purpose
without the express written permission of Rothamsted Research is prohibited.
Printed: Rothamsted, Rothamsted 2008

87/R/B/5 Varieties - W. Barley

Rothamsted Research

Rothamsted Research (1988) *87/R/B/5 Varieties - W. Barley* ; Yields Of The Field Experiments 1987, pp 187 - 188 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/B/5

WINTER BARLEY

VARIETIES

Object: To study the yields of some of the newer winter barley varieties
- Summerdells II.

Sponsors: R. Moffitt, J.F. Jenkyn.

Design: 4 randomised blocks of 11 plots.

Whole plot dimensions: 3.0 x 10.0.

Treatments:

VARIETY	Varieties:
GERBEL	Gerbel
IGRI	Igri
KASKADE	Kaskade
MAGIE	Magie
MG 26+0	Magie with 26 kg extra N applied on 17 Nov, 1986
MG 0+25	Magie with 25 kg extra N applied on 16 Feb, 1987
MG 26+25	Magie with extra N applied on both the above dates
MG S600	Magie with 'Seamac 600' spray
MARINKA	Marinka
PIRATE	Pirate
PLAISANT	Plaisant

NOTES: (1) The extra N for VARIETY MG was applied as urea.
(2) The 'Seamac 600' was applied at 5.6 l in 220 l on 14 Apr, 1987.

Basal applications: Manures: 'Nitram' at 450 kg. Weedkillers: Paraquat at 0.60 kg ion in 200 l. Isoproturon at 2.5 kg with clopyralid at 0.07 kg, bromoxynil at 0.34 kg and mecoprop at 2.5 kg in 200 l. Fungicides: Prochloraz at 0.40 kg and carbendazim at 0.15 kg in 200 l. Triadimenol at 0.12 kg in 200 l. Growth regulators: Mepiquat chloride at 0.61 kg and 2-chloroethylphosphonic acid at 0.31 kg with a wetting agent ('Cittowet' at 0.08 l) in 200 l.

Seed: Varieties sown at 150 kg.

Cultivations, etc.: - Heavy spring-tine cultivated, disced: 18 Aug, 1986. Paraquat applied: 10 Sept. Rotary harrowed, seed sown: 22 Sept. N applied: 19 Mar, 1987. Remaining weedkillers applied: 16 Apr. Prochloraz and carbendazim applied, growth regulators with the wetting agent applied: 27 Apr. Triadimenol applied: 27 May. Combine harvested: 7 Aug. Previous crops: S. barley 1985, w. barley 1986.

NOTE: Leaf samples were taken for disease assessment in June.

87/R/B/5

GRAIN TONNES/HECTARE

***** Tables of means *****

VARIETY	
GERBEL	8.32
IGRI	7.29
KASKADE	7.42
MAGIE	7.95
MG 26+0	8.28
MG 0+25	7.72
MG 26+25	8.58
MG S600	7.77
MARINKA	8.19
PIRATE	7.87
PLAISANT	7.69
Mean	7.92

*** Standard errors of differences of means ***

Table	VARIETY
s.e.d.	0.197

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	30	0.279	3.5
GRAIN MEAN DM%	85.2		
PLOT AREA HARVESTED	0.00204		