

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

ARC, Institute of Arable Crops Research
Rothamsted Experimental Station
Harpenden
Herts
SG8 5LR
UK
The copyright in this document is owned by the Rothamsted Research Ltd
and is a part of the Rothamsted Research Archives. It is made available
on the condition that you will not use it for any other purpose without
the prior written permission of Rothamsted Research Ltd.
Printed: Rothamsted, Bedfordshire
Rothamsted 2008

Field Beans

Rothamsted Research

Rothamsted Research (1988) *Field Beans* ; Yields Of The Field Experiments 1987, pp 192 - 195 -
DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/BE/1

WINTER BEANS

CONTROL OF CHOCOLATE SPOT AND RUST

Object: To compare maneb plus mancozeb with benomyl plus chlorothalonil for the control of chocolate spot (*Botrytis* spp.) and rust (*Uromyces viciae-fabae*) on w. beans sown at two densities - Great Harpenden I.

Sponsors: J. McEwen, D.P. Yeoman.

Design: 2 randomised blocks of 18 plots.

Whole plot dimensions: 6.0 x 10.0.

Treatments: All combinations of:-

1. SEEDRATE Seeds sown per square metre:
 12
 36
2. CS FUNG Fungicides applied to control chocolate spot until first rust pustules seen:

 NONE None
 BEN+CHL Benomyl at 0.50 kg plus chlorothalonil at 1.0 kg on 18 June, 1987
 MAN+MANC Maneb plus mancozeb each at 0.80 kg on 18 June
3. RUSTFUNG Fungicides applied to control rust first applied as soon as rust pustules seen:

 NONE None
 BEN+CHL Benomyl at 0.50 kg plus chlorothalonil at 1.0 kg on 9 July, 5 Aug
 MAN+MANC Maneb plus mancozeb each at 0.80 kg on 9 July, 5 Aug

NOTES: (1) All spray treatments were applied in 200 l.
(2) All benomyl plus chlorothalonil treatments had a wetting agent ('Agral' at 0.06 l) added.

Basal applications: Manures: Chalk at 5.0 t. Weedkillers: Paraquat at 0.80 kg ion in 500 l. Simazine at 1.2 kg with propyzamide at 0.85 kg in 500 l. Insecticide: Deltamethrin at 0.0079 kg in 200 l on two occasions. Desiccant: Diquat at 0.60 kg ion and a wetting agent ('Agral' at 0.3 l) in 300 l.

Seed: Bourdon, dressed with thiram and thiabendazole.

Cultivations, etc.: - Heavy spring-tine cultivated: 10 Sept, 1986. Chalk applied: 24 Sept. Paraquat applied: 6 Nov. Seed broadcast and ploughed in: 12 Nov. Simazine and propyzamide applied: 5 Jan, 1987. Insecticide applied: 22 Apr, 27 May. Desiccant with wetting agent applied: 21 Sept. Combine harvested: 25 Sept. Previous crops: W. wheat 1985 and 1986.

NOTE: Establishment counts were made in autumn, disease assessments were made in July and August and components of yield were measured at maturity.

87/R/BE/1

GRAIN TONNES/HECTARE

***** Tables of means *****

CS FUNG SEEDRATE	NONE	BEN+CHL	MAN+MANC	Mean
12	3.12	2.78	3.40	3.10
36	5.19	5.27	5.11	5.19
Mean	4.15	4.02	4.25	4.14

RUSTFUNG SEEDRATE	NONE	BEN+CHL	MAN+MANC	Mean
12	2.88	3.21	3.20	3.10
36	4.63	5.25	5.68	5.19
Mean	3.76	4.23	4.44	4.14

RUSTFUNG CS FUNG	NONE	BEN+CHL	MAN+MANC	Mean
NONE	3.76	4.25	4.45	4.15
BEN+CHL	3.68	3.90	4.48	4.02
MAN+MANC	3.83	4.54	4.39	4.25
Mean	3.76	4.23	4.44	4.14

SEEDRATE	CS FUNG	RUSTFUNG	NONE	BEN+CHL	MAN+MANC
12	NONE		2.89	3.17	3.29
	BEN+CHL		2.78	2.76	2.80
	MAN+MANC		2.96	3.72	3.51
36	NONE		4.62	5.33	5.61
	BEN+CHL		4.59	5.05	6.17
	MAN+MANC		4.69	5.37	5.26

*** Standard errors of differences of means ***

Table	SEEDRATE	CS FUNG	RUSTFUNG	SEEDRATE CS FUNG
s.e.d.	0.156	0.191	0.191	0.269

Table	SEEDRATE RUSTFUNG	CS FUNG RUSTFUNG	SEEDRATE CS FUNG RUSTFUNG
s.e.d.	0.269	0.330	0.467

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	17	0.467	11.3

GRAIN MEAN DM% 78.0

PLOT AREA HARVESTED 0.00310

87/R/BE/4

SPRING BEANS

VARIETIES, ROW SPACING AND PLANT HEALTH

Object: To compare four varieties of spring-sown field beans at two row spacings and two amounts of pest and disease control - Pastures.

Sponsors: J. McEwen, D.P. Yeoman.

Design: 3 randomised blocks of 16 plots.

Whole plot dimensions: 6.0 x 10.0.

Treatments: All combinations of:-

1. VARIETY Varieties sown at 60 seeds per square metre:
ALFRED
MINDEN
TICOL
TROY
2. ROW SPAC Spacing between rows (cm):
12
48
3. PATHCONT Pest and pathogen control:
STANDARD None
ENHANCED Deltamethrin at 0.0079 kg in 500 l on 22 Apr, 1987,
and in 200 l on 27 May
Chlorothalonil at 1.0 kg with benomy1 at 0.50 kg and
a wetting agent ('Agral' at 0.06 l) in 200 l on
18 June
Maneb at 0.80 kg and mancozeb at 0.80 kg in 200 l on
9 July

Basal applications: Weedkillers: Trietazine at 1.2 kg and simazine at 0.17 kg in 500 l.

Cultivations, etc.:- Ploughed: 27 Oct, 1986. Spring-tine cultivated, rotary harrowed, seed sown: 17 Mar, 1987. Weedkillers applied: 30 Mar. Combine harvested: 21 Sept. Previous crops: W. wheat 1985 and 1986.

NOTE: Establishment counts were made. Disease assessments were made in July and August. At maturity, plant heights, lodging and components of yield were assessed.

87/R/BE/4

GRAIN TONNES/HECTARE

***** Tables of means *****

ROWSPACE	12	48	Mean
VARIETY			
ALFRED	4.88	4.87	4.88
MINDEN	4.47	4.36	4.42
TICOL	3.79	3.68	3.73
TROY	3.95	3.71	3.83
Mean	4.28	4.16	4.22

PATHCONT	STANDARD	ENHANCED	Mean
VARIETY			
ALFRED	3.82	5.94	4.88
MINDEN	3.88	4.95	4.42
TICOL	2.90	4.57	3.73
TROY	3.47	4.20	3.83
Mean	3.52	4.91	4.22

PATHCONT	STANDARD	ENHANCED	Mean
ROWSPACE			
12	3.53	5.03	4.28
48	3.51	4.80	4.16
Mean	3.52	4.91	4.22

	ROWSPACE	12	48		48
VARIETY	PATHCONT	STANDARD	ENHANCED	STANDARD	ENHANCED
ALFRED		3.86	5.91	3.78	5.96
MINDEN		3.98	4.96	3.78	4.93
TICOL		2.87	4.72	2.94	4.42
TROY		3.39	4.51	3.54	3.88

*** Standard errors of differences of means ***

Table	VARIETY	ROWSPACE	PATHCONT	VARIETY ROWSPACE
s.e.d.	0.123	0.087	0.087	0.174

Table	VARIETY PATHCONT	ROWSPACE PATHCONT	VARIETY ROWSPACE PATHCONT
s.e.d.	0.174	0.123	0.246

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	30	0.302	7.2
GRAIN MEAN DM%	76.4		
PLOT AREA HARVESTED	0.00315		