

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1987

[Full Table of Content](#)

ARC, Institute of Arable Crops Research  
Rothamsted Experimental Station  
Harpenden  
Herts  
AL5 2JQ  
UK  
The copyright in this document is held by the Rothamsted Research Ltd.  
This document is available on the internet at <https://www.era-doc.org/>  
This document is available on the internet at <https://www.era-doc.org/>  
Printed: Rothamsted, Rothamsted 1988

## 87/R/BE/4 Varieties, Row Spacing and Plant Health - S. Beans

### Rothamsted Research

Rothamsted Research (1988) *87/R/BE/4 Varieties, Row Spacing and Plant Health - S. Beans* ; Yields Of The Field Experiments 1987, pp 194 - 195 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/BE/4

SPRING BEANS

VARIETIES, ROW SPACING AND PLANT HEALTH

Object: To compare four varieties of spring-sown field beans at two row spacings and two amounts of pest and disease control - Pastures.

Sponsors: J. McEwen, D.P. Yeoman.

Design: 3 randomised blocks of 16 plots.

Whole plot dimensions: 6.0 x 10.0.

Treatments: All combinations of:-

1. VARIETY Varieties sown at 60 seeds per square metre:  
ALFRED  
MINDEN  
TICOL  
TROY
2. ROW SPAC Spacing between rows (cm):  
12  
48
3. PATHCONT Pest and pathogen control:  
STANDARD None  
ENHANCED Deltamethrin at 0.0079 kg in 500 l on 22 Apr, 1987,  
and in 200 l on 27 May  
Chlorothalonil at 1.0 kg with benomy1 at 0.50 kg and  
a wetting agent ('Agral' at 0.06 l) in 200 l on  
18 June  
Maneb at 0.80 kg and mancozeb at 0.80 kg in 200 l on  
9 July

Basal applications: Weedkillers: Trietazine at 1.2 kg and simazine at 0.17 kg in 500 l.

Cultivations, etc.:- Ploughed: 27 Oct, 1986. Spring-tine cultivated, rotary harrowed, seed sown: 17 Mar, 1987. Weedkillers applied: 30 Mar. Combine harvested: 21 Sept. Previous crops: W. wheat 1985 and 1986.

NOTE: Establishment counts were made. Disease assessments were made in July and August. At maturity, plant heights, lodging and components of yield were assessed.

87/R/BE/4

GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

ROWSpace	12	48	Mean
VARIETY			
ALFRED	4.88	4.87	4.88
MINDEN	4.47	4.36	4.42
TICOL	3.79	3.68	3.73
TROY	3.95	3.71	3.83
Mean	4.28	4.16	4.22

PATHCONT	STANDARD	ENHANCED	Mean
VARIETY			
ALFRED	3.82	5.94	4.88
MINDEN	3.88	4.95	4.42
TICOL	2.90	4.57	3.73
TROY	3.47	4.20	3.83
Mean	3.52	4.91	4.22

PATHCONT	STANDARD	ENHANCED	Mean
ROWSpace			
12	3.53	5.03	4.28
48	3.51	4.80	4.16
Mean	3.52	4.91	4.22

	ROWSpace	12	48		48
VARIETY	PATHCONT	STANDARD	ENHANCED	STANDARD	ENHANCED
ALFRED		3.86	5.91	3.78	5.96
MINDEN		3.98	4.96	3.78	4.93
TICOL		2.87	4.72	2.94	4.42
TROY		3.39	4.51	3.54	3.88

\*\*\* Standard errors of differences of means \*\*\*

Table	VARIETY	ROWSpace	PATHCONT	VARIETY ROWSpace
s.e.d.	0.123	0.087	0.087	0.174

Table	VARIETY PATHCONT	ROWSpace PATHCONT	VARIETY ROWSpace PATHCONT
s.e.d.	0.174	0.123	0.246

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	30	0.302	7.2
GRAIN MEAN DM%	76.4		
PLOT AREA HARVESTED	0.00315		