

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

ROTHAMSTED RESEARCH
Rothamsted Experimental Station
Harpenden
Herts
AL5 2JQ
UK
TEL: 0456 473900
FAX: 0456 473901
E-MAIL: enquiries@ Rothamsted.ac.uk
WWW: <http:// Rothamsted.ac.uk>
© Rothamsted Research 2008

87/R/LP/4 Desiccants and Fungicides - Lupins

Rothamsted Research

Rothamsted Research (1988) *87/R/LP/4 Desiccants and Fungicides - Lupins* ; Yields Of The Field Experiments 1987, pp 199 - 201 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/LP/4

LUPINS

DESICCANTS AND FUNGICIDES

Object: To study the effects of times of applying desiccants and fungicides on senescence, grain quality and yield of lupins (*Lupinus albus*) - Long Hoos III 1.

Sponsors: H.L. Jones, J. Lacey.

Design: 2 randomised blocks of 30 plots.

Whole plot dimensions: 2.4 x 5.0.

Treatments: All combinations of:-

- | | |
|--------------|--|
| 1. DESICCANT | Desiccants: |
| DIQUAT | Diquat at 0.15 kg ion |
| MET+FEN | Metoxuron at 2.0 kg plus fentin hydroxide at 0.20 kg |
| GLYPHOS | Glyphosate at 1.08 kg |
| NACL | Sodium chloride at 6.25 kg |
| 2. FUNGICIDE | Fungicide: |
| NONE | None |
| PROPICON | Propiconazole at 0.125 kg |
| 3. APP TIME | Times of applying desiccants and fungicide: |
| EARLY | Fungicide on 25 Sept 1987, desiccants 12 Oct |
| MIDDLE | Fungicide on 25 Sept, desiccants 19 Oct |
| LATE | Fungicide on 25 Sept, desiccants 28 Oct |

plus five extra treatments not given desiccants:

- | | |
|----------|---|
| FUNGIC X | Fungicides applied on 12 Oct: |
| NONE | None (duplicated) |
| IPRODION | Iprodione at 1.0 kg |
| PROPICON | Propiconazole at 0.125 kg |
| PROP+DIQ | Propiconazole at 0.125 kg + diquat at 0.60 kg ion |
| VINCLOZ | Vinclozalin at 0.375 kg |

- NOTES: (1) APP TIME for desiccants was chosen for the three stages EARLY-all leaves below pods fallen, MIDDLE-pods fully developed, LATE-no leaves remaining on plant.
(2) All spray treatments were applied in 220 l.
(3) The crop was netted from early Apr to early June.

Basal applications: Weedkillers: Glyphosate at 1.44 kg in 220 l. Monolinuron at 0.46 kg with paraquat at 0.33 kg ion in 220 l. Metamitron at 2.8 kg in 220 l. Insecticides: Cypermethrin at 0.025 kg in 220 l. Pirimicarb at 0.14 kg in 220 l applied with the fungicide. Fungicide: Benomyl at 0.55 kg.

Seed: Vladimir, dressed with thiram, inoculated with Rhizobium, sown at 240 kg.

87/R/LP/4

Cultivations, etc.:— Glyphosate applied: 1 Oct, 1986. Ploughed: 17-29 Oct. Seed sown, harrowed, rolled: 31 Mar, 1987. Monolinuron and paraquat applied: 2 Apr. Cypermethrin applied: 8 May. Metamitron applied: 18 June. Benomyl and pirimicarb applied: 13 July. Combine harvested: 17 Nov. Previous crops: S. beans 1985, fallow 1986.

NOTE: Plant populations were measured and disease assessments made during the season. After desiccation, changes in dry matter were measured and fungal development on the pods assessed. Components of yield were measured. Seed microflora, germinability and discolouration of the seed coat were also assessed.

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

FUNGCIDE	NONE	PROPICON	Mean			
DESICCNT						
DIQUAT	1.63	1.89	1.76			
MET+FEN	1.79	1.46	1.62			
GLYPHOS	1.59	1.71	1.65			
NACL	1.81	1.80	1.80			
Mean	1.71	1.71	1.71			
APP TIME	EARLY	MIDDLE	LATE	Mean		
DESICCNT						
DIQUAT	1.59	1.92	1.77	1.76		
MET+FEN	1.34	1.81	1.73	1.62		
GLYPHOS	1.93	1.68	1.35	1.65		
NACL	1.86	2.01	1.55	1.80		
Mean	1.68	1.85	1.60	1.71		
APP TIME	EARLY	MIDDLE	LATE	Mean		
FUNGCIDE						
NONE	1.73	1.90	1.49	1.71		
PROPICON	1.63	1.80	1.71	1.71		
Mean	1.68	1.85	1.60	1.71		
DESICCNT	APP TIME	EARLY	MIDDLE	LATE		
DIQUAT	FUNGCIDE					
NONE		1.60	1.78	1.52		
PROPICON		1.57	2.07	2.03		
MET+FEN	FUNGCIDE					
NONE		1.61	2.08	1.69		
PROPICON		1.07	1.53	1.76		
GLYPHOS	FUNGCIDE					
NONE		1.91	1.59	1.28		
PROPICON		1.94	1.77	1.43		
NACL	FUNGCIDE					
NONE		1.79	2.17	1.47		
PROPICON		1.93	1.84	1.62		
FUNGIC X	NONE	IPRODION	PROPICON	PROP+DIQ	VINCLOZ	Mean
	1.78	1.54	1.71	1.41	1.84	1.68
Grand mean	1.70					

87/R/LP/4

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

*** Standard errors of differences of means ***

Table	DESICCNT	FUNGCIDE	APP TIME	DESICCNT FUNGCIDE
s.e.d.	0.162	0.114	0.140	0.228

Table	DESICCNT APP TIME	FUNGCIDE APP TIME	DESICCNT FUNGCIDE APP TIME	FUNGC X
s.e.d.	0.280	0.198	0.396	0.396

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	30	0.396	23.2

GRAIN MEAN DM% 63.3

PLOT AREA HARVESTED 0.00072