

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

ARC, Institute of Arable Crops Research
Rothamsted Experimental Station
Harpenden
Herts
SG8 5LR
UK
The copyright in this document is owned by the Rothamsted Research and
is licensed to the user of this document for personal use only. No part of
this document may be reproduced, stored in a retrieval system, or
transmitted in any form or by any means, electronic, mechanical, or
otherwise, without the prior written permission of Rothamsted Research.
Printed: Northampton, 1988
Rothamsted 1988

87/R/LP/4 Desiccants and Fungicides - Lupins

Rothamsted Research

Rothamsted Research (1988) *87/R/LP/4 Desiccants and Fungicides - Lupins* ; Yields Of The Field Experiments 1987, pp 199 - 201 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/LP/4

LUPINS

DESICCANTS AND FUNGICIDES

Object: To study the effects of times of applying desiccants and fungicides on senescence, grain quality and yield of lupins (*Lupinus albus*) - Long Hoos III 1.

Sponsors: H.L. Jones, J. Lacey.

Design: 2 randomised blocks of 30 plots.

Whole plot dimensions: 2.4 x 5.0.

Treatments: All combinations of:-

- | | |
|--------------|--|
| 1. DESICCANT | Desiccants: |
| DIQUAT | Diquat at 0.15 kg ion |
| MET+FEN | Metoxuron at 2.0 kg plus fentin hydroxide at 0.20 kg |
| GLYPHOS | Glyphosate at 1.08 kg |
| NACL | Sodium chloride at 6.25 kg |
| 2. FUNGICIDE | Fungicide: |
| NONE | None |
| PROPICON | Propiconazole at 0.125 kg |
| 3. APP TIME | Times of applying desiccants and fungicide: |
| EARLY | Fungicide on 25 Sept 1987, desiccants 12 Oct |
| MIDDLE | Fungicide on 25 Sept, desiccants 19 Oct |
| LATE | Fungicide on 25 Sept, desiccants 28 Oct |

plus five extra treatments not given desiccants:

- | | |
|----------|---|
| FUNGIC X | Fungicides applied on 12 Oct: |
| NONE | None (duplicated) |
| IPRODION | Iprodione at 1.0 kg |
| PROPICON | Propiconazole at 0.125 kg |
| PROP+DIQ | Propiconazole at 0.125 kg + diquat at 0.60 kg ion |
| VINCLOZ | Vinclozalin at 0.375 kg |

- NOTES: (1) APP TIME for desiccants was chosen for the three stages EARLY-all leaves below pods fallen, MIDDLE-pods fully developed, LATE-no leaves remaining on plant.
(2) All spray treatments were applied in 220 l.
(3) The crop was netted from early Apr to early June.

Basal applications: Weedkillers: Glyphosate at 1.44 kg in 220 l. Monolinuron at 0.46 kg with paraquat at 0.33 kg ion in 220 l. Metamitron at 2.8 kg in 220 l. Insecticides: Cypermethrin at 0.025 kg in 220 l. Pirimicarb at 0.14 kg in 220 l applied with the fungicide. Fungicide: Benomyl at 0.55 kg.

Seed: Vladimir, dressed with thiram, inoculated with Rhizobium, sown at 240 kg.

87/R/LP/4

Cultivations, etc.:- Glyphosate applied: 1 Oct, 1986. Ploughed: 17-29 Oct. Seed sown, harrowed, rolled: 31 Mar, 1987. Monolinuron and paraquat applied: 2 Apr. Cypermethrin applied: 8 May. Metamitron applied: 18 June. Benomyl and pirimicarb applied: 13 July. Combine harvested: 17 Nov. Previous crops: S. beans 1985, fallow 1986.

NOTE: Plant populations were measured and disease assessments made during the season. After desiccation, changes in dry matter were measured and fungal development on the pods assessed. Components of yield were measured. Seed microflora, germinability and discolouration of the seed coat were also assessed.

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

| | | | | | | |
|------------|----------|----------|----------|----------|---------|------|
| FUNGCIDE | NONE | PROPICON | Mean | | | |
| DESICCNT | | | | | | |
| DIQUAT | 1.63 | 1.89 | 1.76 | | | |
| MET+FEN | 1.79 | 1.46 | 1.62 | | | |
| GLYPHOS | 1.59 | 1.71 | 1.65 | | | |
| NACL | 1.81 | 1.80 | 1.80 | | | |
| Mean | 1.71 | 1.71 | 1.71 | | | |
| APP TIME | EARLY | MIDDLE | LATE | Mean | | |
| DESICCNT | | | | | | |
| DIQUAT | 1.59 | 1.92 | 1.77 | 1.76 | | |
| MET+FEN | 1.34 | 1.81 | 1.73 | 1.62 | | |
| GLYPHOS | 1.93 | 1.68 | 1.35 | 1.65 | | |
| NACL | 1.86 | 2.01 | 1.55 | 1.80 | | |
| Mean | 1.68 | 1.85 | 1.60 | 1.71 | | |
| APP TIME | EARLY | MIDDLE | LATE | Mean | | |
| FUNGCIDE | | | | | | |
| NONE | 1.73 | 1.90 | 1.49 | 1.71 | | |
| PROPICON | 1.63 | 1.80 | 1.71 | 1.71 | | |
| Mean | 1.68 | 1.85 | 1.60 | 1.71 | | |
| DESICCNT | APP TIME | EARLY | MIDDLE | LATE | | |
| DIQUAT | FUNGCIDE | | | | | |
| NONE | | 1.60 | 1.78 | 1.52 | | |
| PROPICON | | 1.57 | 2.07 | 2.03 | | |
| MET+FEN | NONE | 1.61 | 2.08 | 1.69 | | |
| PROPICON | | 1.07 | 1.53 | 1.76 | | |
| GLYPHOS | NONE | 1.91 | 1.59 | 1.28 | | |
| PROPICON | | 1.94 | 1.77 | 1.43 | | |
| NACL | NONE | 1.79 | 2.17 | 1.47 | | |
| PROPICON | | 1.93 | 1.84 | 1.62 | | |
| FUNGIC X | NONE | IPRODION | PROPICON | PROP+DIQ | VINCLOZ | Mean |
| | 1.78 | 1.54 | 1.71 | 1.41 | 1.84 | 1.68 |
| Grand mean | 1.70 | | | | | |

87/R/LP/4

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

*** Standard errors of differences of means ***

| | | | | |
|--------|----------|----------|----------|----------------------|
| Table | DESICCNT | FUNGCIDE | APP TIME | DESICCNT FUNGCIDE |
| s.e.d. | 0.162 | 0.114 | 0.140 | 0.228 |

| | | | | |
|--------|----------------------|----------------------|----------------------------------|---------|
| Table | DESICCNT APP TIME | FUNGCIDE APP TIME | DESICCNT FUNGCIDE APP TIME | FUNGC X |
| s.e.d. | 0.280 | 0.198 | 0.396 | 0.396 |

***** Stratum standard errors and coefficients of variation *****

| | | | |
|----------|------|-------|------|
| Stratum | d.f. | s.e. | cv% |
| BLOCK.WP | 30 | 0.396 | 23.2 |

GRAIN MEAN DM% 63.3

PLOT AREA HARVESTED 0.00072