

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

ARC, Institute of Arable Crops Research
Rothamsted Experimental Station
Harpenden
Herts
SG8 5LR
UK

The copyright in this document is owned by the Rothamsted Research and is licensed to the National Institute for Research in Food Safety and Food Quality (NIFFSQ) for the purpose of making available to the public the full text of this document. All other rights reserved.

Printed: 2016/06/06
Rothamsted 2016

87/R/RA/11 Oversowing in Wheat - W. Oilseed Rape

Rothamsted Research

Rothamsted Research (1988) *87/R/RA/11 Oversowing in Wheat - W. Oilseed Rape* ; Yields Of The Field Experiments 1987, pp 235 - 236 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/RA/11

WINTER OILSEED RAPE

OVERSOWING IN WHEAT

Object: To study the establishment of rape after wheat by oversowing into the wheat - Great Knott III.

Sponsors: R.J. Darby, D.P. Yeoman.

Design: 3 randomised blocks of 3 plots split into 3 sub plots.

Whole plot dimensions: 15.0 x 16.0.

Treatments: All combinations of:-

Whole plots

- | | |
|-----------|---|
| 1. SOWING | Methods of sowing and straw disposal: |
| OVERS BA | Oversown on 20 Aug, 1986, straw baled on 21 Aug |
| OVERS CH | Oversown on 20 Aug, 1986, straw chopped and spread on 21 Aug |
| CONVEN S | Straw baled on 21 Aug, conventionally sown into conventionally prepared seedbed |

Sub plots

- | | |
|-----------|---|
| 2. N TIME | Timing of nitrogen as 'Nitram'. (Total spring N same for all treatments): |
| SN FN MN | 50 kg N to seedbed, 50 kg N on 19 Feb, 1987 and 150 kg N on 19 Mar |
| - FN - | 200 kg N on 19 Feb |
| - FN MN | 50 kg N on 19 Feb and 150 kg N on 19 Mar |

- NOTES: (1) Oversowing was done into standing wheat. The wheat was harvested later that day.
(2) SOWING CONVEN S plots were ploughed and rolled on 22 Aug, rotary harrowed on 31 Aug.
(3) SOWING OVERS BA and OVERS CH plots were sprayed with fluazifop-P-butyl at 0.19 kg with a wetting agent ('Agral' at 0.38 l) in 380 l on 26 Sept.
(4) One whole plot CONVEN S was missing. An estimated value was used in the analysis.

Basal applications: Weedkillers: Clopyralid and propyzamide (as 'Matrikerb' at 1.6 kg) in 500 l. Desiccant: Diquat at 0.60 kg ion with a wetting agent ('Agral' at 0.50 l) in 500 l.

Seed: Ariana, dressed gamma HCH, thiram and fenpropimorph, sown at 8.0 kg.

Cultivations, etc.:- Weedkillers applied: 20 Nov, 1986. Desiccant with wetting agent applied: 31 July, 1987. Combine harvested: 6 Aug. Previous crops: W. wheat 1985 and 1986.

NOTE: Plant counts were made in autumn and again in mid-March.

87/R/RA/11

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

	N TIME	SN FN MN	- FN -	- FN MN	Mean
SOWING					
OVERS BA		3.61	3.51	3.38	3.50
OVERS CH		3.47	2.86	2.66	3.00
CONVEN S		3.51	3.28	3.22	3.34
Mean		3.53	3.22	3.09	3.28

*** Standard errors of differences of means ***

Table	N TIME	SOWING*
s.e.d.	0.184	0.319

* Within same level of SOWING only

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP.SP	10	0.391	11.9

GRAIN MEAN DM% 75.2

SUB PLOT AREA HARVESTED 0.00368