

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

ARC, Institute of Arable Crops Research
Rothamsted Experimental Station
Harpenden
Herts
SG8 5LR
UK
The copyright in this document is owned by the Rothamsted Research Ltd
and is a part of the Rothamsted Research Archives. It is made available
online as a service to the public. It is not to be used for any other
purpose without the prior written permission of Rothamsted Research Ltd.
Printed: 2018/08/08
Rothamsted 2018

87/R/SU/3 Pests and Pathogens - Sunflowers

Rothamsted Research

Rothamsted Research (1988) *87/R/SU/3 Pests and Pathogens - Sunflowers* ; Yields Of The Field Experiments 1987, pp 242 - 243 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/SU/3

SUNFLOWERS

PESTS AND PATHOGENS

Object: To study the separate and combined effects of pest and pathogen control on sunflowers - Garden Plot 2.

Sponsors: A.W. Ferguson, C.J. Rawlinson.

Design: 4 randomised blocks of 4 plots.

Whole plot dimensions: 2.13 x 10.1.

Treatments: All combinations of:-

1. PESTCONT Sprays to control pests:

NONE	None
FULL	Full: Pirimicarb at 0.14 kg on 25 July, 1987, 5, 19 Aug, 3, 17 Sept. Malathion at 1.3 l on 4, 11, 19 Aug

2. PATHCONT Sprays to control pathogens:

NONE	None
FULL	Full: Carbendazim at 0.25 kg with vinclozolin at 0.50 kg on 25 July, 5, 19 Aug, 3, 17 Sept, 1 Oct

- NOTES: (1) Pesticides were applied in the same 220 l when applied on the same day otherwise in 220 l each. Fungicides were always applied together in 220 l.
(2) The crop was netted against bird damage from late Aug to harvest.
(3) A planned test of varieties was thwarted by bird damage.

Basal applications : Manures: 'Nitro-Chalk' at 225 kg. Weedkillers: Glyphosate at 1.4 kg in 220 l. Trifluralin at 1.1 kg in 220 l. Linuron at 0.50 kg in 220 l. Desiccant: Diquat at 0.60 kg ion in 220 l.

Seed: Asmer 9, sown at 90,000 seeds per hectare.

Cultivations, etc.:- Glyphosate applied: 1 Oct, 1986. Ploughed: 15 Oct. N applied: 21 Apr, 1987. Trifluralin applied, rotary cultivated: 23 Apr. Seed sown, rolled: 24 Apr. Linuron applied: 29 Apr. Diquat applied: 19 Oct. Harvested by hand: 23 Oct (threshed by stationary combine harvester). Previous crops: Mixed legumes 1985, s. barley 1986.

- NOTES: (1) Aphids and pollen beetles were counted weekly in late summer.
(2) Growth stages and diseases were observed on six occasions during the season.

87/R/SU/3

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

PATHCONT	NONE	FULL	Mean
PESTCONT			
NONE	1.98	2.06	2.02
FULL	1.84	2.05	1.95
Mean	1.91	2.05	1.98

*** Standard errors of differences of means ***

Table	PESTCONT	PATHCONT	PESTCONT PATHCONT
s.e.d.	0.110	0.110	0.156

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	9	0.221	11.1

GRAIN MEAN DM% 60.0

PLOT AREA HARVESTED 0.00215