Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1987



Full Table of Content

87/R/SU/4 Fungicide and Botrytis - Sunflowers

Rothamsted Research

Rothamsted Research (1988) 87/R/SU/4 Fungicide and Botrytis - Sunflowers; Yields Of The Field Experiments 1987, pp 244 - 246 - **DOI:** https://doi.org/10.23637/ERADOC-1-37

87/R/SU/4

SUNFLOWERS

FUNGICIDE AND BOTRYTIS

Object: To study the effects of five times and two methods of applying a mixture of fungicides on the control of Botrytis head infection and on the yield of sunflowers - Annables and Long Hoos III 3.

Sponsor: C.J. Rawlinson.

Design: 3 randomised blocks of 12 plots on each site.

Whole plot dimensions: Annables: 4.2 x 7.0. Long Hoos: 2.7 x 8.0.

Treatments: All combinations of:-

1.	SPRAYDTE	Dates	of	applying	vinclozolin	at	0.50	kg	and
		cal	rbei	ndazim at	0.25 kg:				

	Annables	Long Hoos
1	25 July, 1987	21 Aug
2	25 July and 5 Aug	21 Aug and 4 Sept
3	25 July, 5 and 19 Aug	21 Aug, 4 and 17 Sept
4	25 July, 5, 19 Aug and 3 Sept	21 Aug, 4, 17 Sept and 1 Oct (duplicated)
5	25 July, 5, 19 Aug, 3 and 16 Sept	d

2. SPRAYMET Method of spraying:

ELECTRO Electrostatic sprayer in 5.3 1 (Annables only)
CNVNTIAL Conventional hydraulic sprayer in 220 1 (duplicated on Long Hoos)

plus one extra treatment

EXTRA

NONE None (duplicated)

NOTE: On Long Hoos the crop was netted from early June to harvest.

Basal applications:

Annables: Manures: 'Nitram' at 170 kg. Weedkiller: Trifluralin at 1.1 kg in 220 l. Desiccant: Diquat at 0.60 kg ion in 220 l. Long Hoos: Manures: 'Nitro-Chalk' at 255 kg. Weedkillers: Trifluralin at 1.1 kg in 220 l. Linuron at 0.50 kg in 220 l. Desiccant: Diquat at 0.60 kg ion in 220 l.

Seed: Annables: Asmer 3 sown at 90,000 seeds per hectare.

Long Hoos: Asmer 3 failed and site was resown with EX10, both at
90,000 seeds per hectare.

87/R/SU/4

Cultivations, etc .:-

Annables: Ploughed: Autumn 1986, date not recorded. Rotary cultivated, 'Nitram' applied, trifluralin applied, rotary cultivated, seed sown: 15 Apr, 1987. Desiccant applied: 23 Sept. Combine harvested: 29 Sept. Previous crops: W. wheat 1985 and 1986.

Long Hoos: Ploughed: 29 Dec, 1986. 'Nitro-Chalk' applied: 21 Apr, 1987. Spring-tine cultivated, trifluralin applied, rotary cultivated, Asmer 3 sown and rolled: 24 Apr. Linuron applied: 29 Apr. EX10 sown: 28 May. Desiccant applied: 15 Oct. Harvested by hand: 23 Oct (threshed by stationary combine harvester). Previous crops: Fallow 1985, lupins 1986.

NOTE: Botrytis in heads and stems and growth stages were assessed on both sites. More detailed observations on crop growth were made on Annables.

ANNABLES

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

**** Tables of means ****

SPRAYMET	ELECTRO	CNVNTIAL	Mean
SPRAYDTE	0.70	0.74	0.73
1	0.73	0.74	
2	0.85	0.79	0.82
3	0.77	0.76	0.77
4	0.77	0.79	0.78
5	0.72	0.72	0.72
Mean	0.77	0.76	0.76

EXTRA NONE

0.84

Grand mean

0.78

*** Standard errors of differences of means ***

Table	SPRAYDTE	SPRAYMET	SPRAYDTE SPRAYMET
s.e.d.	0.128	0.081	0.181

SED for comparing NONE with any item in SPRAYMET.SPRAYDTE table is 0.156

**** Stratum standard errors and coefficients of variation ****

Stratum d.f. s.e. cv%
BLOCK.WP 23 0.221 28.5

GRAIN MEAN DM% 69.2

PLOT AREA HARVESTED 0.00223

87/R/SU/4

LONG HOOS III

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

SPRAYDTE NONE 1 2 3 4 5 Mean 1.02 0.99 1.11 1.10 1.04 0.91 1.03

*** Standard errors of differences of means ***

Table

SPRAYDTE

s.e.d.

0.093

**** Stratum standard errors and coefficients of variation ****

Stratum

d.f.

s.e.

CV%

BLOCK . WP

27

0.161

15.6

GRAIN MEAN DM% 48.2

PLOT AREA HARVESTED 0.00216