

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

ARC, Institute of Arable Crops Research
Rothamsted Experimental Station
Harpenden
Herts
SG8 5LR
UK
The copyright in this document is owned by the Rothamsted Research Ltd
and is a trademark of Rothamsted Research Ltd. All rights reserved.
This document is available on the internet at <http://www.era.doc>
Printed: 2010/08/08
Rothamsted 2008

87/R/BR/1 Anti-feedants - Broccoli

Rothamsted Research

Rothamsted Research (1988) *87/R/BR/1 Anti-feedants - Broccoli* ; Yields Of The Field Experiments 1987, pp 252 - 253 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/BR/1

BROCCOLI

ANTI-FEEDANTS

Object: To study the effects of anti-feedants on insect pests and on the yield of broccoli - Garden Plot 7.

Sponsors: D.C. Griffiths, L.E. Smart.

Design: 5 randomised blocks of 4 plots.

Whole plot dimensions: 2.4 x 5.0.

Treatments:

CHEMICAL Chemical foliar sprays:

NONE	None
AJUGA EX	Ajuga extract at 0.15 kg
DELTAMET	Deltamethrin at 0.01 kg
PIPERINE	Piperine at 0.20 kg

NOTES: (1) Treatments were applied on 7 July, 1987, 24 July, 30 July, 11 Aug, 20 Aug, 4 Sept, 11 Sept, 21 Sept and 1 Oct.
(2) They were applied by electrostatic sprayer in 10 l.

Basal applications: Manures: Chalk at 2.9 t. 'Nitro-Chalk' at 390 kg.
Weedkillers: Glyphosate at 1.4 kg in 220 l, bentazone at 0.80 kg, dichlorprop at 1.1 kg, MCPA at 0.64 kg in 220 l applied with the fungicide. Trifluralin at 1.1 kg in 220 l. Tebutam at 2.9 kg in 220 l. Fungicide: Tridemorph at 0.52 kg.

Seed: Sown at 140,000 seeds per hectare.

Cultivations, etc.:- Chalk applied: 29 Sept, 1986. Glyphosate applied: 1 Oct. Ploughed: 22 Dec. Bentazone, dichlorprop, MCPA and tridemorph applied: 6 May, 1987. Deep-tine cultivated twice, trifluralin applied, rotary cultivated, seed sown, rolled, tebutam applied: 7 May. 'Nitro-Chalk' applied: 11 May. Seed resown: 22 June. Harvested by hand: 4 Nov. Previous crops: Potatoes 1985, fallow 1986.

NOTES: (1) The first sowing failed and was resown.
(2) Flea beetle, aphid and caterpillar damage were assessed at intervals during the season.

87/R/BR/1

FRESH WEIGHT TONNES/HECTARE

***** Tables of means *****

CHEMICAL	NONE	AJUGA EX	DELTAMET	PIPERINE	Mean
	33.8	31.0	34.0	27.8	31.7

*** Standard errors of differences of means ***

Table	CHEMICAL
s.e.d.	2.63

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	12	4.16	13.2
PLOT AREA HARVESTED	0.00048		