

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

ARC, Institute of Arable Crops Research
Rothamsted Experimental Station
Harpenden
Herts
SG8 5AA
ENGLAND
UK

The copyright in this document is owned by the Rothamsted Research Ltd. All rights are reserved and no part of this document may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of Rothamsted Research.

Printed: Northampton, ENGLAND 2008

Mixed Crops

Rothamsted Research

Rothamsted Research (1988) *Mixed Crops* ; Yields Of The Field Experiments 1987, pp 268 - 278 -
DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/M/1 and 87/W/M/1

MIXED 1

INPUTS FOR WINTER CEREALS

Object: To compare amounts of disease and the yield of triticale with those of w. wheat, w. barley and w. rye on two contrasted sites each given contrasted amounts of agrochemicals - Rothamsted Summerdells I (R), Woburn Great Hill II (W).

Sponsors: R.J. Gutteridge, D. Hornby, R.D. Prew (R), P.R. Scott, W. Hollins, R.L. Gregory (P.B.I., Cambridge).

Design: 3 randomised blocks of 10 plots.

Whole plot dimensions: 3.0 x 10.0 (R), 4.0 x 10.0 (W).

Treatments: All combinations of :-

1. CROP VAR	Crop and variety:	(R)	(W)
B PANDA	W. barley, Panda sown at	230 kg	230 kg
R DOMINT	W. rye, Dominant sown at	170 kg	170 kg
T LASKO	W. triticale, Lasko sown at	170 kg	160 kg
T CWT	W. triticale, CWT/1977/290 sown at	180 kg	170 kg
W AVALON	W. wheat, Avalon sown at	190 kg	190 kg

2. INPUT Inputs of agrochemicals, in addition to basals:

LARGE	(R): Manures: N at 40 kg: 11 Feb, 1987, and at 160 kg: 2 Apr, both as 'Nitro-Chalk'. Fungicides: Prochloraz at 0.40 kg, carbendazim at 0.15 kg, tridemorph at 0.52 kg in 220 l: 8 May. Carbendazim at 0.25 kg, maneb at 1.6 kg with propiconazole at 0.12 kg in 220 l: 2 July. Growth regulators: Mepiquat chloride at 0.61 kg with 2-chloroethylphosphonic acid at 0.31 kg in 220 l to barley, chlormequat at 1.1 kg in 220 l to wheat and triticale: 8 May.
	(W): Manures: N at 40 kg: 13 Feb, 1987 and at 160 kg: 31 Mar, both as 'Nitram'. Fungicides: Prochloraz at 0.40 kg, carbendazim at 0.15 kg, tridemorph at 0.52 kg in 240 l: 21 Apr. Propiconazole at 0.12 kg, tridemorph at 0.52 kg in 200 l: 27 May. Propiconazole at 0.12 kg, carbendazim at 0.25 kg in 200 l: 29 June. Growth regulators: Mepiquat chloride at 0.53 kg with 2-chloroethylphosphonic acid at 0.27 kg in 200 l, to barley and triticale: 7 May.
SMALL	(R) Manures: 120 kg N as 'Nitro-Chalk': 2 Apr, 1987. (W) Manures: 160 kg N as 'Nitram': 31 Mar.

87/R/M/1 and 87/W/M/1

Basal applications:

Summerdells (R): Manures: Chalk at 5.0 t. Weedkillers: Paraquat at 0.60 kg ion in 200 l. Methabenzthiazuron at 1.6 kg in 200 l. Isoproturon at 2.5 kg with bromoxynil and ioxynil (as 'Deloxil' at 2.0 l) in 380 l to barley only. Diclofop-methyl at 1.1 kg with bromoxynil and ioxynil (as 'Deloxil' at 2.0 l) in 380 l to rye, triticale and wheat.

Great Hill II (W): Weedkillers: Bromoxynil and ioxynil (as 'Deloxil' at 2.0 l) in 240 l. Fluroxypyr at 0.20 kg in 400 l to barley and wheat.

Cultivations, etc.:-

Summerdells (R): Heavy spring-tine cultivated and disced: 19 Aug, 1986. Chalk applied: 4 Sept. Paraquat applied: 11 Sept. Spring-tine cultivated, rotary harrowed, seed sown, harrowed: 24 Sept. Rolled: 27 Sept. Methabenzthiazuron applied: 30 Sept. Isoproturon, bromoxynil and ioxynil applied to barley, diclofop-methyl, bromoxynil and ioxynil applied to rye, triticale and wheat: 17 Apr, 1987. Combine harvested barley: 7 Aug, rye, triticale and wheat: 1 Sept. Previous crops: W. wheat 1985, w. barley 1986.

Great Hill II (W): Ploughed, rolled: 20 Sept, 1986. Rotary harrowed with crumbler attached, seed sown: 25 Sept. Bromoxynil and ioxynil applied: 17 Apr, 1987. Fluroxypyr applied to barley and wheat: 23 Apr. Combine harvested barley: 5 Aug, rye, triticale and wheat: 18 Aug. Previous crops: Lucerne 1985, w. wheat 1986.

- NOTES: (1) Soil samples were taken for take-all bioassay before sowing and after harvest.
(2) Assessments were made of foot and root rots and foliar diseases during the season.

87/R/M/1

GRAIN TONNES/HECTARE

***** Tables of means *****

INPUT	LARGE	SMALL	Mean
CROP VAR			
B PANDA	6.59	6.46	6.53
R DOMINT	6.54	6.04	6.29
T LASKO	5.74	4.62	5.18
T CWT	6.02	5.81	5.91
W AVALON	6.66	6.17	6.42
Mean	6.31	5.82	6.06

*** Standard errors of differences of means ***

Table	CROP VAR	INPUT	CROP VAR INPUT
s.e.d.	0.270	0.170	0.381

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	18	0.467	7.7

GRAIN MEAN DM% 85.0

PLOT AREA HARVESTED 0.00274

87/W/M/1

GRAIN TONNES/HECTARE

***** Tables of means *****

INPUT	LARGE	SMALL	Mean
CROP VAR			
B PANDA	6.56	6.16	6.36
R DOMINT	6.39	5.66	6.02
T LASKO	4.42	3.79	4.11
T CWT	4.57	4.70	4.64
W AVALON	4.72	3.70	4.21
Mean	5.33	4.80	5.07

*** Standard errors of differences of means ***

Table	CROP VAR	INPUT	CROP VAR INPUT
s.e.d.	0.396	0.251	0.560

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	18	0.686	13.5

GRAIN MEAN DM% 80.1

PLOT AREA HARVESTED 0.00275

87/R/M/2

MIXED 2

FACTORS AFFECTING EYESPOT

Object: To study eyespot (*Pseudocercospora herpotrichoides*) development after inoculation with different pathotypes in relation to host crop and seed rate - White Horse II.

Sponsor: A. Goulds.

Design: 2 blocks of 12 plots split into 3.

Whole plot dimensions: 3.0 x 37.0.

Treatments: All combinations of:-

Whole plots

1. W CEREAL Winter cereals sown on 30 September, 1986:
 BARLEY Winter barley cv. Opera
 WHEAT Winter wheat cv. Avalon
2. SEEDRATE Seed rates (seeds per square metre):
 NORMAL Normal - 300 barley, 400 wheat
 HALF N Half normal - 150 barley, 200 wheat
3. INOCULUM Inoculation with different eyespot pathogen types:
 NONE None
 RYE INOC Rye type
 WHE INOC Wheat type

Sub plots

4. FUNGTIME Times of applying prochloraz at 0.40 kg and carbendazim at 0.15 kg in 220 l:
 NONE None
 EARLY Sprayed at growth stage 30/31 on 23 Apr, 1987
 LATE Sprayed at growth stage, 33/37 wheat, 41/49 barley on 19 May

NOTES: (1) One additional sub-plot in each whole plot was systematically arranged for sampling, yields not taken.
(2) Strains of wheat and rye type inoculum were colonised on oat seed and broadcast within two weeks of emergence.

Basal applications: Manures: 'Nitram' at 350 kg. Weedkillers: Isoproturon at 2.5 kg with clopyralid at 0.07 kg and bromoxynil at 0.34 kg and mecoprop at 2.5 kg in 200 l. Fungicides: Propiconazole at 0.25 kg with tridemorph at 0.19 kg in 200 l.

Cultivations, etc.:- Heavy spring-tine cultivated twice, disced: 29 Sept, 1986. Rotary harrowed, seed sown: 30 Sept. Weedkillers applied: 15 Apr, 1987. N applied: 17 Apr. Basal fungicides applied: 29 June. Combine harvested: 7 Aug (barley), 1 Sept (wheat). Previous crops: W oats 1985, potatoes 1986.

87/R/M/2

NOTE: Eyespot was assessed on plants at fortnightly intervals from December to G.S. 30 and weekly thereafter.

GRAIN TONNES/HECTARE

***** Tables of means *****

SEEDRATE	NORMAL	HALF N	Mean	
W CEREAL				
BARLEY	7.74	7.67	7.71	
WHEAT	7.98	7.85	7.92	
Mean	7.86	7.76	7.81	
INOCULUM	NONE	RYE INOC	WHE INOC	Mean
W CEREAL				
BARLEY	7.97	7.74	7.41	7.71
WHEAT	7.94	8.03	7.77	7.92
Mean	7.95	7.89	7.59	7.81
INOCULUM	NONE	RYE INOC	WHE INOC	Mean
SEEDRATE				
NORMAL	8.09	7.91	7.57	7.86
HALF N	7.81	7.87	7.61	7.76
Mean	7.95	7.89	7.59	7.81
FUNGTIME	NONE	EARLY	LATE	Mean
W CEREAL				
BARLEY	7.18	8.06	7.88	7.71
WHEAT	7.56	7.95	8.23	7.92
Mean	7.37	8.00	8.05	7.81
FUNGTIME	NONE	EARLY	LATE	Mean
SEEDRATE				
NORMAL	7.32	8.08	8.18	7.86
HALF N	7.43	7.93	7.92	7.76
Mean	7.37	8.00	8.05	7.81
FUNGTIME	NONE	EARLY	LATE	Mean
INOCULUM				
NONE	7.47	8.22	8.17	7.95
RYE INOC	7.72	7.78	8.16	7.89
WHE INOC	6.92	8.02	7.82	7.59
Mean	7.37	8.00	8.05	7.81
W CEREAL	INOCULUM	NONE	RYE INOC	WHE INOC
BARLEY	SEEDRATE			
NORMAL		8.15	7.83	7.24
HALF N		7.78	7.65	7.58
WHEAT	NORMAL	8.04	7.98	7.91
	HALF N	7.85	8.08	7.63

87/R/M/2

GRAIN TONNES/HECTARE

***** Tables of means *****

	FUNGTIME	NONE	EARLY	LATE
W CEREAL	SEEDRATE			
	BARLEY			
	NORMAL	7.03	8.17	8.02
	HALF N	7.33	7.95	7.73
WHEAT	NORMAL	7.60	7.98	8.35
	HALF N	7.53	7.92	8.12
	FUNGTIME	NONE	EARLY	LATE
W CEREAL	INOCULUM			
	BARLEY			
	NONE	7.54	8.18	8.17
	RYE INOC	7.45	7.99	7.79
WHEAT	WHE INOC	6.55	8.02	7.66
	NONE	7.40	8.25	8.18
	RYE INOC	8.00	7.57	8.53
	WHE INOC	7.29	8.03	7.99
	FUNGTIME	NONE	EARLY	LATE
SEEDRATE	INOCULUM			
	NORMAL			
	NONE	7.56	8.36	8.36
	RYE INOC	7.77	7.74	8.22
HALF N	WHE INOC	6.62	8.13	7.97
	NONE	7.39	8.07	7.99
	RYE INOC	7.68	7.82	8.10
	WHE INOC	7.23	7.91	7.68
	FUNGTIME	NONE	EARLY	LATE
W CEREAL	INOCULUM			
	BARLEY			
	NONE	7.74	8.39	8.33
	RYE INOC	7.52	8.03	7.96
	WHE INOC	5.84	8.11	7.78
HALF N	NONE	7.34	7.97	8.02
	RYE INOC	7.39	7.95	7.63
WHEAT	WHE INOC	7.27	7.93	7.54
	NONE	7.38	8.34	8.39
	RYE INOC	8.02	7.45	8.49
	WHE INOC	7.40	8.16	8.17
HALF N	NONE	7.43	8.17	7.96
	RYE INOC	7.98	7.69	8.58
	WHE INOC	7.19	7.89	7.81

87/R/M/2

GRAIN TONNES/HECTARE

*** Standard errors of differences of means ***

Table	W CEREAL	SEEDRATE	INOCULUM	FUNGTIME
s.e.d.	0.093	0.093	0.114	0.114
Table	W CEREAL SEEDRATE	W CEREAL INOCULUM	SEEDRATE INOCULUM	W CEREAL FUNGTIME
s.e.d.	0.132	0.162	0.162	0.162
Table	SEEDRATE FUNGTIME	INOCULUM FUNGTIME	W CEREAL SEEDRATE INOCULUM	W CEREAL SEEDRATE FUNGTIME
s.e.d.	0.162	0.198	0.229	0.228
Table	W CEREAL INOCULUM FUNGTIME	SEEDRATE INOCULUM FUNGTIME	W CEREAL SEEDRATE INOCULUM FUNGTIME	
s.e.d.	0.280	0.280	0.396	

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	11	0.229	2.9
BLOCK.WP.SP	24	0.396	5.1

GRAIN MEAN DM% 84.3

SUB PLOT AREA HARVESTED 0.00235

87/W/M/2

MIXED 2

COMPARISON OF COMBINES

Object: To evaluate the suitability of two combines for plot work in respect of purity of sample and accuracy when working on slopes - Great Hill III.

Sponsor: R. Moffitt.

Design: A systematic split-plot design, 56 whole plots arranged as shown below:

R	W	R	W	R	W	R	W	Top of slope
W	W	W	W	W	W	W	W	
W	W	W	W	W	W	W	W	
R	R	R	R	R	R	R	R	
W	W	W	W	W	W	W	W	
W	W	W	W	W	W	W	W	
R*	W	R	W	R	W	R	W	Bottom of slope

* Combines started here (after harvesting a dummy wheat plot downhill), worked up the column of plots then down the next column etc.

R = Rye W = wheat

- NOTES: (1) Each whole plot was systematically divided to compare the two combine harvesters.
(2) There were 10 m headlands between contiguous rye and wheat plots. These were removed before combining the plots. There were 1 m paths between contiguous wheat plots.

Whole plot dimensions: 8.0 x 11.0.

Treatments:

Whole plots

- CROP Crop and variety:
 WHEAT W. wheat, Avalon
 RYE W. rye, Dominant
- DIRECTN Combine direction in relation to slope:
 UP Up slope
 DOWN Down slope
- ORDER Order of combining:
 BEGIN First plot in column
 STRAIGHT Central plots in column
 END Last plot in column

87/W/M/2

Sub plots

4. COMBINE Combine type:

CLAYSON	Clayson 1530
DEUTZ-F	Deutz-Fahr 660

Basal applications: Manures: FYM at 100 t. (0:18:36) at 700 kg, 'Nitram' at 470 kg. Weedkillers: Bromoxynil and ioxynil (as 'Deloxil' at 2.0 l) in 240 l. Fungicide: Triadimenol at 0.062 kg in 200 l.

Seed: W. rye: Dominant, sown at 170 kg.
W. wheat: Avalon, sown at 160 kg.

Cultivations, etc.: - FYM applied: 3 July, 1986. Ploughed: 9-11, July. Rolled: 22 July. PK applied: 16 Sept. Rotary harrowed with crumbler attached, sown: 7 Oct. Rolled: 8 Oct. N applied: 15 Apr, 1987. Weedkillers applied: 17 June. Fungicide applied: 1 July. Combine harvested: 15 Sept. Previous crops: Lucerne 1985 and 1986.

87/W/M/2

GRAIN TONNES/HECTARE

***** Tables of means *****

COMBINE	CLAYSON	DEUTZ-F					
	5.63	5.78					
	DIRECTN	UP				DOWN	
CROP	ORDER	BEGIN	STRAIGHT	END	BEGIN	STRAIGHT	END
WHEAT			6.40		5.49	6.58	6.67
RYE		3.94	3.65	4.25		3.93	
CROP	DIRECTN	ORDER	COMBINE	CLAYSON	DEUTZ-F		
WHEAT	UP	STRAIGHT		6.24	6.55		
	DOWN	BEGIN		5.57	5.41		
		STRAIGHT		6.46	6.70		
		END		6.55	6.78		
RYE	UP	BEGIN		4.49	3.39		
		STRAIGHT		3.76	3.54		
		END		3.80	4.70		
	DOWN	STRAIGHT		3.78	4.09		

Grand mean 5.70

*** Standard errors of differences of means ***

Table	COMBINE	CROP DIRECTN ORDER	CROP DIRECTN ORDER COMBINE	
s.e.d.	0.066	0.242	0.299	min.rep
		0.191	0.236	max-min
		0.121	0.149	max.rep
Except when comparing means with the same level(s) of				
CROP.DIRECTN.ORDER			0.249	min.rep
			0.197	max-min
			0.124	max.rep

max.rep CROP WHEAT and ORDER STRAIGHT

min.rep any of the remainder

max-min CROP WHEAT and ORDER STRAIGHT v any of the remainder

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
WP	48	0.342	6.0
WP.SP	48	0.352	6.2

GRAIN MEAN DM% 82.0

PLOT AREA HARVESTED	CLAYSON	DEUTZ-F
	0.00302	0.00220