

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

ROTHAMSTED RESEARCH
ROTHAMSTED EXPERIMENTAL STATION
ROTHAMSTED
Herts
SG2 0AT
UK
TEL: 0454 464400
FAX: 0454 464401
WWW: www.rothamsted.ac.uk
© Rothamsted Research 1988

87/R/M/2 Factors Affecting Eyespot - W. Wheat, Barley - Mixed Crops

Rothamsted Research

Rothamsted Research (1988) *87/R/M/2 Factors Affecting Eyespot - W. Wheat, Barley - Mixed Crops* ;
Yields Of The Field Experiments 1987, pp 272 - 275 - DOI:

<https://doi.org/10.23637/ERADOC-1-37>

87/R/M/2

MIXED 2

FACTORS AFFECTING EYESPOT

Object: To study eyespot (*Pseudocercospora herpotrichoides*) development after inoculation with different pathotypes in relation to host crop and seed rate - White Horse II.

Sponsor: A. Goulds.

Design: 2 blocks of 12 plots split into 3.

Whole plot dimensions: 3.0 x 37.0.

Treatments: All combinations of:-

Whole plots

- | | |
|-------------|--|
| 1. W CEREAL | Winter cereals sown on 30 September, 1986: |
| BARLEY | Winter barley cv. Opera |
| WHEAT | Winter wheat cv. Avalon |
| 2. SEEDRATE | Seed rates (seeds per square metre): |
| NORMAL | Normal - 300 barley, 400 wheat |
| HALF N | Half normal - 150 barley, 200 wheat |
| 3. INOCULUM | Inoculation with different eyespot pathogen types: |
| NONE | None |
| RYE INOC | Rye type |
| WHE INOC | Wheat type |

Sub plots

- | | |
|-------------|--|
| 4. FUNGTIME | Times of applying prochloraz at 0.40 kg and carbendazim at 0.15 kg in 220 l: |
| NONE | None |
| EARLY | Sprayed at growth stage 30/31 on 23 Apr, 1987 |
| LATE | Sprayed at growth stage, 33/37 wheat, 41/49 barley on 19 May |

NOTES: (1) One additional sub-plot in each whole plot was systematically arranged for sampling, yields not taken.
(2) Strains of wheat and rye type inoculum were colonised on oat seed and broadcast within two weeks of emergence.

Basal applications: Manures: 'Nitram' at 350 kg. Weedkillers: Isoproturon at 2.5 kg with clopyralid at 0.07 kg and bromoxynil at 0.34 kg and mecoprop at 2.5 kg in 200 l. Fungicides: Propiconazole at 0.25 kg with tridemorph at 0.19 kg in 200 l.

Cultivations, etc.:- Heavy spring-tine cultivated twice, disced: 29 Sept, 1986. Rotary harrowed, seed sown: 30 Sept. Weedkillers applied: 15 Apr, 1987. N applied: 17 Apr. Basal fungicides applied: 29 June. Combine harvested: 7 Aug (barley), 1 Sept (wheat). Previous crops: W oats 1985, potatoes 1986.

87/R/M/2

NOTE: Eyespot was assessed on plants at fortnightly intervals from December to G.S. 30 and weekly thereafter.

GRAIN TONNES/HECTARE

***** Tables of means *****

SEEDRATE	NORMAL	HALF N	Mean	
W CEREAL				
BARLEY	7.74	7.67	7.71	
WHEAT	7.98	7.85	7.92	
Mean	7.86	7.76	7.81	
INOCULUM	NONE	RYE INOC	WHE INOC	Mean
W CEREAL				
BARLEY	7.97	7.74	7.41	7.71
WHEAT	7.94	8.03	7.77	7.92
Mean	7.95	7.89	7.59	7.81
INOCULUM	NONE	RYE INOC	WHE INOC	Mean
SEEDRATE				
NORMAL	8.09	7.91	7.57	7.86
HALF N	7.81	7.87	7.61	7.76
Mean	7.95	7.89	7.59	7.81
FUNGTIME	NONE	EARLY	LATE	Mean
W CEREAL				
BARLEY	7.18	8.06	7.88	7.71
WHEAT	7.56	7.95	8.23	7.92
Mean	7.37	8.00	8.05	7.81
FUNGTIME	NONE	EARLY	LATE	Mean
SEEDRATE				
NORMAL	7.32	8.08	8.18	7.86
HALF N	7.43	7.93	7.92	7.76
Mean	7.37	8.00	8.05	7.81
FUNGTIME	NONE	EARLY	LATE	Mean
INOCULUM				
NONE	7.47	8.22	8.17	7.95
RYE INOC	7.72	7.78	8.16	7.89
WHE INOC	6.92	8.02	7.82	7.59
Mean	7.37	8.00	8.05	7.81
W CEREAL	INOCULUM	NONE	RYE INOC	WHE INOC
BARLEY	SEEDRATE			
NORMAL		8.15	7.83	7.24
HALF N		7.78	7.65	7.58
WHEAT	NORMAL	8.04	7.98	7.91
	HALF N	7.85	8.08	7.63

87/R/M/2

GRAIN TONNES/HECTARE

***** Tables of means *****

	FUNGTIME	NONE	EARLY	LATE
W CEREAL	SEEDRATE			
	BARLEY			
	NORMAL	7.03	8.17	8.02
	HALF N	7.33	7.95	7.73
WHEAT	NORMAL	7.60	7.98	8.35
	HALF N	7.53	7.92	8.12
	FUNGTIME	NONE	EARLY	LATE
W CEREAL	INOCULUM			
	BARLEY			
	NONE	7.54	8.18	8.17
	RYE INOC	7.45	7.99	7.79
WHEAT	WHE INOC	6.55	8.02	7.66
	NONE	7.40	8.25	8.18
	RYE INOC	8.00	7.57	8.53
	WHE INOC	7.29	8.03	7.99
	FUNGTIME	NONE	EARLY	LATE
SEEDRATE	INOCULUM			
	NORMAL			
	NONE	7.56	8.36	8.36
	RYE INOC	7.77	7.74	8.22
HALF N	WHE INOC	6.62	8.13	7.97
	NONE	7.39	8.07	7.99
	RYE INOC	7.68	7.82	8.10
	WHE INOC	7.23	7.91	7.68
	FUNGTIME	NONE	EARLY	LATE
W CEREAL	INOCULUM			
	BARLEY			
	NONE	7.74	8.39	8.33
	RYE INOC	7.52	8.03	7.96
	WHE INOC	5.84	8.11	7.78
HALF N	NONE	7.34	7.97	8.02
	RYE INOC	7.39	7.95	7.63
WHEAT	WHE INOC	7.27	7.93	7.54
	NONE	7.38	8.34	8.39
	RYE INOC	8.02	7.45	8.49
	WHE INOC	7.40	8.16	8.17
HALF N	NONE	7.43	8.17	7.96
	RYE INOC	7.98	7.69	8.58
	WHE INOC	7.19	7.89	7.81

87/R/M/2

GRAIN TONNES/HECTARE

*** Standard errors of differences of means ***

Table	W CEREAL	SEEDRATE	INOCULUM	FUNGTIME
s.e.d.	0.093	0.093	0.114	0.114
Table	W CEREAL SEEDRATE	W CEREAL INOCULUM	SEEDRATE INOCULUM	W CEREAL FUNGTIME
s.e.d.	0.132	0.162	0.162	0.162
Table	SEEDRATE FUNGTIME	INOCULUM FUNGTIME	W CEREAL SEEDRATE INOCULUM	W CEREAL SEEDRATE FUNGTIME
s.e.d.	0.162	0.198	0.229	0.228
Table	W CEREAL INOCULUM FUNGTIME	SEEDRATE INOCULUM FUNGTIME	W CEREAL SEEDRATE INOCULUM FUNGTIME	
s.e.d.	0.280	0.280	0.396	

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	11	0.229	2.9
BLOCK.WP.SP	24	0.396	5.1

GRAIN MEAN DM% 84.3

SUB PLOT AREA HARVESTED 0.00235