

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

ARC, Institute of Arable Crops Research
Rothamsted Experimental Station
Harpenden
Herts
SG8 5LR
United Kingdom
ARC 1988
The copyright in this document is held by the Rothamsted Research Institute. It is published as part of the Rothamsted Research Reports. The copyright in this document is held by the Rothamsted Research Institute. It is published as part of the Rothamsted Research Reports.

87/R/RN/17 Rates of P and K to the Subsoil - S. Beans, W. Wheat, Potatoes, S. Barley

Rothamsted Research

Rothamsted Research (1988) *87/R/RN/17 Rates of P and K to the Subsoil - S. Beans, W. Wheat, Potatoes, S. Barley*; Yields Of The Field Experiments 1987, pp 70 - 76 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/RN/17

RATES OF P AND K TO THE SUBSOIL

Object: To study the effects of a range of rates and frequencies of application of P and K to the subsoil, singly and together, on the yields and nutrient uptakes of a rotation of crops - Meadow.

Sponsors: J. McEwen, A.E. Johnston.

The seventh year, potatoes, s. barley, s. beans, w. wheat.

For previous years see 81-86/R/RN/17.

Design: 4 series (for crops) each of 40 plots.

Whole plot dimensions: 3.0 x 14.0.

Treatments to each series:

TREATMNT Extra P and K and primary cultivation tool in autumn 1980 only, except on A plots, treatments repeated annually, and F plots treatments repeated four yearly:

| | P205(kg) | K20(kg) | Tool | |
|----------|----------|----------------|-------------------|---------------|
| - - - | 0 | 0 | Plough | (duplicated) |
| P6 K6 T | 1000 | 500 to topsoil | " | (") |
| - - S | 0 | 0 | Wye double-digger | (triplicated) |
| - - SA | 0 | 0 | " " | (duplicated) |
| - - SF | 0 | 0 | " " | " |
| P2 - SA | 63 | 0 to subsoil | " " | " |
| P3 - SF | 125 | 0 " " | " " | " |
| P4 - S | 250 | 0 " " | " " | " |
| P5 - S | 500 | 0 " " | " " | " |
| P5 - SF | 500 | 0 " " | " " | " |
| P6 - S | 1000 | 0 " " | " " | " |
| - K2 SA | 0 | 31 " " | " " | " |
| - K3 SF | 0 | 63 " " | " " | " |
| - K4 S | 0 | 125 " " | " " | " |
| - K5 S | 0 | 250 " " | " " | " |
| - K5 SF | 0 | 250 " " | " " | " |
| - K6 S | 0 | 350 " " | " " | " |
| P1 K1 SA | 31 | 16 " " | " " | " |
| P1 K3 SA | 31 | 63 " " | " " | " |
| P2 K2 SA | 63 | 31 " " | " " | " |
| P3 K1 SA | 125 | 16 " " | " " | " |
| P3 K3 SA | 125 | 63 " " | " " | " |
| P3 K4 SF | 125 | 125 " " | " " | " |
| P4 K3 SF | 250 | 63 " " | " " | " |
| P4 K4 S | 250 | 125 " " | " " | " |
| P4 K5 S | 250 | 250 " " | " " | " |
| P4 K5 SF | 250 | 250 " " | " " | " |
| P4 K6 S | 250 | 350 " " | " " | " |
| P5 K4 S | 500 | 125 " " | " " | " |
| P5 K4 SF | 500 | 125 " " | " " | " |
| P5 K5 S | 500 | 250 " " | " " | " |
| P5 K6 S | 500 | 350 " " | " " | " |
| P6 K4 S | 1000 | 125 " " | " " | " |
| P6 K5 S | 1000 | 250 " " | " " | " |
| P6 K6 S | 1000 | 350 " " | " " | " |

87/R/RN/17

- NOTES: (1) Subsoiling was done with the Wye double-digger which turns a furrow with a conventional plough share, to a depth of 23 cm, and at the same time rotary cultivates the bottom of the adjacent furrow to a further depth of 15 cm. When applying P and K this was distributed ahead of the rotary cultivator.
- (2) The topsoil PK dressing was equally divided before and after ploughing.
- (3) All plots were conventionally ploughed each autumn unless the Wye double-digging treatment was due.
- (4) The rate of 350 kg K20 applied was in error for 500 kg K20.

Standard applications:

- Potatoes: Manures: (10:10:15+4.5 Mg) at 1960 kg. Weedkiller: Linuron at 1.6 kg in 500 l. Fungicides: Mancozeb at 1.4 kg in 200 l on four occasions, applied with the pirimicarb on the second occasion. Fentin hydroxide at 0.28 kg in 200 l on two occasions. Insecticide: Pirimicarb at 0.14 kg. Desiccant: Diquat at 0.80 kg in 300 l.
- S. barley: Manures: (20:10:10) at 630 kg. Weedkillers: Clopyralid at 0.07 kg, bromoxynil at 0.34 kg and mecoprop at 2.5 kg in 200 l. Fungicide: Tridemorph at 0.52 kg in 200 l.
- S. beans: Weedkillers: Trietazine at 1.2 kg and simazine at 0.17 kg in 500 l. Insecticide: Phorate at 4.5 kg.
- W. wheat: Manures: (0:18:36) at 350 kg. 'Nitram' at 590 kg. Weedkillers: Clopyralid at 0.07 kg and bromoxynil at 0.34 kg with mecoprop at 2.5 kg 200 l. Fungicide: Tridemorph at 0.52 kg in 200 l.

Seed: Potatoes: Pentland Crown.

- S. barley: Klaxon, dressed triadimenol and fuberidazole, sown at 160 kg.
- S. beans: Minden, sown at 260 kg.
- W. wheat: Avalon, sown at 200 kg.

Cultivations, etc.:-

All crops: Treatments applied by double-digger: 29 Oct, 1986 and 30 Oct. Ploughed: 31 Oct.

- Potatoes: Heavy spring-tine cultivated: 20 Feb, 1987. NPK+Mg applied: 14 Apr. Heavy spring-tine cultivated: 15 Apr. Rotary harrowed, potatoes planted: 16 Apr. Rotary ridged: 27 Apr. Linuron applied: 8 May. Mancozeb applied: 24 June, 8 July, 28 July, 10 Aug. Insecticide applied: 8 July. Fentin hydroxide applied: 28 Aug, 9 Sept. Desiccant applied: 21 Sept. Haulm mechanically destroyed: 3 Oct. Lifted: 19 Oct.
- S. barley: Heavy spring-tine cultivated: 20 Feb, 1987. NPK applied, spring-tine cultivated: 16 Mar. Rotary harrowed, seed sown: 17 Mar. Weedkillers applied: 6 May. Fungicide applied: 24 June. Combine harvested: 20 Aug.
- S. beans: Heavy spring-tine cultivated: 20 Feb, 1987. Phorate applied, spring-tine cultivated: 16 Mar. Rotary harrowed, seed sown: 18 Mar. Weedkillers applied: 30 Mar. Combine harvested: 18 Sept.
- W. wheat: PK applied, rotary harrowed, seed sown: 7 Nov, 1986. N applied: 17 Apr, 1987. Weedkillers applied: 8 May. Fungicide applied: 24 June. Combine harvested: 31 Aug.

87/R/RN/17

SERIES III POTATOES

TOTAL TUBERS TONNES/HECTARE

***** Tables of means *****

| TREATMNT | |
|----------|------|
| - - - | 64.6 |
| P6 K6 T | 60.0 |
| - - S | 61.7 |
| - - SA | 62.2 |
| - - SF | 63.6 |
| P2 - SA | 62.4 |
| P3 - SF | 71.5 |
| P4 - S | 50.7 |
| P5 - S | 65.4 |
| P5 - SF | 66.9 |
| P6 - S | 68.1 |
| - K2 SA | 62.6 |
| - K3 SF | 65.0 |
| - K4 S | 68.1 |
| - K5 S | 70.1 |
| - K5 SF | 66.6 |
| - K6 S | 70.8 |
| P1 K1 SA | 61.8 |
| P1 K3 SA | 67.6 |
| P2 K2 SA | 69.5 |
| P3 K1 SA | 67.9 |
| P3 K3 SA | 65.7 |
| P3 K4 SF | 71.8 |
| P4 K3 SF | 63.0 |
| P4 K4 S | 62.3 |
| P4 K5 S | 65.0 |
| P4 K5 SF | 68.8 |
| P4 K6 S | 68.1 |
| P5 K4 S | 68.7 |
| P5 K4 SF | 67.8 |
| P5 K5 S | 70.5 |
| P5 K6 S | 64.6 |
| P6 K4 S | 67.0 |
| P6 K5 S | 67.7 |
| P6 K6 S | 69.2 |
| Mean | 65.4 |

*** Standard errors of differences of means ***

| Table | TREATMNT |
|--------|--------------|
| s.e.d. | 7.14 min.rep |
| | 5.83 max-min |

***** Stratum standard errors and coefficients of variation *****

| Stratum | d.f. | s.e. | cv% |
|---------|------|------|-----|
| WP | 5 | 5.05 | 7.7 |

87/R/RN/17

SERIES III POTATOES

PERCENTAGE WARE 3.81 CM (1.5 INCH) RIDDLE

***** Tables of means *****

| TREATMNT | |
|----------|------|
| - - - | 97.6 |
| P6 K6 T | 98.5 |
| - - S | 98.2 |
| - - SA | 98.5 |
| - - SF | 97.6 |
| P2 - SA | 97.8 |
| P3 - SF | 98.8 |
| P4 - S | 97.7 |
| P5 - S | 98.5 |
| P5 - SF | 98.0 |
| P6 - S | 98.3 |
| - K2 SA | 98.5 |
| - K3 SF | 98.1 |
| - K4 S | 97.7 |
| - K5 S | 97.7 |
| - K5 SF | 98.3 |
| - K6 S | 98.1 |
| P1 K1 SA | 96.7 |
| P1 K3 SA | 98.4 |
| P2 K2 SA | 98.2 |
| P3 K1 SA | 99.1 |
| P3 K3 SA | 97.3 |
| P3 K4 SF | 98.8 |
| P4 K3 SF | 98.2 |
| P4 K4 S | 97.9 |
| P4 K5 S | 98.3 |
| P4 K5 SF | 97.9 |
| P4 K6 S | 98.5 |
| P5 K4 S | 97.5 |
| P5 K4 SF | 98.1 |
| P5 K5 S | 99.3 |
| P5 K6 S | 96.9 |
| P6 K4 S | 98.4 |
| P6 K5 S | 98.1 |
| P6 K6 S | 99.3 |
| Mean | 98.1 |

PLOT AREA HARVESTED 0.00210

* SEDs apply only to - - -, P6 K6 T, - - S and - - SA

| | TREATMNT |
|---------|--------------------------|
| max-min | - - S v any of remainder |
| min.rep | any of remainder |

87/R/RN/17

SERIES IV WINTER BARLEY

GRAIN TONNES/HECTARE

***** Tables of means *****

| TREATMNT | |
|----------|------|
| - - - | 7.02 |
| P6 K6 T | 6.92 |
| - - S | 6.95 |
| - - SA | 7.22 |
| - - SF | 6.73 |
| P2 - SA | 7.42 |
| P3 - SF | 7.12 |
| P4 - S | 6.99 |
| P5 - S | 6.98 |
| P5 - SF | 7.23 |
| P6 - S | 6.50 |
| - K2 SA | 7.28 |
| - K3 SF | 7.10 |
| - K4 S | 7.21 |
| - K5 S | 7.37 |
| - K5 SF | 7.05 |
| - K6 S | 6.92 |
| P1 K1 SA | 7.10 |
| P1 K3 SA | 6.82 |
| P2 K2 SA | 6.90 |
| P3 K1 SA | 6.82 |
| P3 K3 SA | 7.40 |
| P3 K4 SF | 7.18 |
| P4 K3 SF | 7.10 |
| P4 K4 S | 7.01 |
| P4 K5 S | 7.30 |
| P4 K5 SF | 7.33 |
| P4 K6 S | 7.12 |
| P5 K4 S | 7.09 |
| P5 K4 SF | 6.86 |
| P5 K5 S | 7.08 |
| P5 K6 S | 7.05 |
| P6 K4 S | 6.58 |
| P6 K5 S | 6.36 |
| P6 K6 S | 7.00 |
| Mean | 7.03 |

*** Standard errors of differences of means ***

| Table | TREATMNT |
|--------|--------------------------------|
| s.e.d. | 0.206 min.rep 0.168 max-min |

***** Stratum standard errors and coefficients of variation *****

| Stratum | d.f. | s.e. | cv% |
|----------------|------|---------------------|---------|
| WP | 5 | 0.146 | 2.1 |
| GRAIN MEAN DM% | 86.0 | PLOT AREA HARVESTED | 0.00286 |

87/R/RN/17

SERIES I SPRING BEANS

GRAIN TONNES/HECTARE

***** Tables of means *****

| TREATMNT | |
|----------|------|
| - - - | 4.50 |
| P6 K6 T | 4.31 |
| - - S | 4.51 |
| - - SA | 4.83 |
| - - SF | 4.14 |
| P2 - SA | 4.34 |
| P3 - SF | 4.43 |
| P4 - S | 4.07 |
| P5 - S | 4.14 |
| P5 - SF | 4.97 |
| P6 - S | 4.68 |
| - K2 SA | 4.74 |
| - K3 SF | 4.77 |
| - K4 S | 4.80 |
| - K5 S | 4.41 |
| - K5 SF | 4.81 |
| - K6 S | 4.72 |
| P1 K1 SA | 4.63 |
| P1 K3 SA | 4.53 |
| P2 K2 SA | 4.88 |
| P3 K1 SA | 4.70 |
| P3 K3 SA | 4.01 |
| P3 K4 SF | 4.75 |
| P4 K3 SF | 4.32 |
| P4 K4 S | 3.82 |
| P4 K5 S | 4.50 |
| P4 K5 SF | 4.27 |
| P4 K6 S | 4.63 |
| P5 K4 S | 5.04 |
| P5 K4 SF | 4.93 |
| P5 K5 S | 4.43 |
| P5 K6 S | 4.80 |
| P6 K4 S | 4.55 |
| P6 K5 S | 4.44 |
| P6 K6 S | 4.44 |
| Mean | 4.54 |

*** Standard errors of differences of means ***

| Table | TREATMNT |
|--------|---------------|
| s.e.d. | 0.430 min.rep |
| | 0.351 max-min |

***** Stratum standard errors and coefficients of variation *****

| Stratum | d.f. | s.e. | cv% |
|----------------|------|---------------------|---------|
| WP | 5 | 0.304 | 6.7 |
| GRAIN MEAN DM% | 81.9 | PLOT AREA HARVESTED | 0.00386 |

87/R/RN/17

SERIES II WINTER WHEAT

GRAIN TONNES/HECTARE

***** Tables of means *****

| TREATMNT | |
|----------|------|
| - - - | 6.69 |
| P6 K6 T | 7.41 |
| - - S | 6.62 |
| - - SA | 6.90 |
| - - SF | 5.93 |
| P2 - SA | 7.10 |
| P3 - SF | 7.79 |
| P4 - S | 7.15 |
| P5 - S | 7.28 |
| P5 - SF | 6.99 |
| P6 - S | 6.85 |
| - K2 SA | 6.81 |
| - K3 SF | 5.53 |
| - K4 S | 7.35 |
| - K5 S | 6.74 |
| - K5 SF | 7.08 |
| - K6 S | 5.17 |
| P1 K1 SA | 7.58 |
| P1 K3 SA | 7.24 |
| P2 K2 SA | 7.49 |
| P3 K1 SA | 7.56 |
| P3 K3 SA | 7.27 |
| P3 K4 SF | 8.14 |
| P4 K3 SF | 7.24 |
| P4 K4 S | 7.29 |
| P4 K5 S | 6.65 |
| P4 K5 SF | 7.01 |
| P4 K6 S | 7.88 |
| P5 K4 S | 6.33 |
| P5 K4 SF | 7.31 |
| P5 K5 S | 6.29 |
| P5 K6 S | 6.82 |
| P6 K4 S | 6.90 |
| P6 K5 S | 6.63 |
| P6 K6 S | 7.10 |
| Mean | 6.96 |

*** Standard errors of differences of means ***

| Table | TREATMNT |
|--------|---------------|
| s.e.d. | 0.624 min.rep |
| | 0.510 max-min |

***** Stratum standard errors and coefficients of variation *****

| Stratum | d.f. | s.e. | cv% |
|----------------|------|---------------------|---------|
| WP | 5 | 0.441 | 6.3 |
| GRAIN MEAN DM% | 83.0 | PLOT AREA HARVESTED | 0.00274 |