

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

# Yields of the Field Experiments 1987

[Full Table of Content](#)

ARC, Institute of Arable Crops Research  
Rothamsted Experimental Station  
Harpenden  
Herts  
SG8 2EQ  
UK  
The copyright in this document is owned by the Rothamsted Research Ltd  
and is a trademark of Rothamsted Research Ltd. All rights reserved. No  
part of this document may be reproduced, stored in a retrieval system,  
transmitted, or otherwise disseminated in any form or by any means  
without the prior written permission of Rothamsted Research Ltd.  
Printed: Rothamsted, Bedfordshire  
Rothamsted 2008

## Table of Means

### Rothamsted Research

Rothamsted Research (1988) *Table of Means* ; Yields Of The Field Experiments 1987, pp 8 - 8 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

### Harvest areas for cereals

On most of those cereal experiments at Rothamsted and Woburn which are harvested by combine the 'blank-row' technique is used to distinguish the areas taken for yield from the discard areas. For example when seed is drilled in rows 7 in. (18 cm) apart appropriate coulters are prevented from sowing and 8 or 16 rows are left for yield according to the cutter-bar width of the combine to be used. If the row-spacing is other than 7 in. a similar arrangement is used but with a different number of rows.

The ends of plots are separated from each other or from headlands by 3 ft (91 cm) fallow paths made after the crop has established.

The 'Area harvested' in the 'Yields', when the blank-row technique is used, is the product:-

number of rows harvested x distance between rows x length of rows.

A series of experiments at Rothamsted showed that on average the yield of 16 rows (50 ft (15 m) long) was 7.8% greater with blank rows than without. (Experimental Husbandry 23 pp 16-20 (1972)).

If no rows are left blank and the plot is wider than the combine harvester so that discards are left uncut, the 'Area harvested' is the product:-

width of cutter bar x length of rows.

If the plot is narrower than the combine so that the whole area between paths is cut, the 'Area harvested' is the product:-

number of rows x distance between rows x length of rows.

We do not apply the adjustment used by some workers who take the harvested areas as width x length where each is measured to the centre of 'paths' up to a maximum of 18 in. (46 cm).

### Tables of means

Tables of means are presented directly from computer output. Both factor and level names are presented in upper case characters. Vertical and horizontal lines are omitted e.g.:-

FACTOR C	LEVEL C1		LEVEL C2		LEVEL C3	
FACTOR B	LEVEL B1	LEVEL B2	LEVEL B1	LEVEL B2	LEVEL B1	LEVEL B2
FACTOR A						
LEVEL A1	*	*	*	*	*	*
LEVEL A2	*	*	*	*	*	*

### Standard errors

- NOTES: (1) This report gives standard errors of differences, not of means.  
 (2) Annotations (e.g. \* min rep, max-min, max rep) to S.E.Ds are only explained the first time they occur in any experiment.