

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

ARC, Institute of Arable Crops Research
Rothamsted Experimental Station
Harpenden
Herts
AL5 2JQ
UK
ARC 1988
The copyright in this document is held by the Rothamsted Research Institute. It is published as part of the Rothamsted Research Reports series. All rights reserved. No part of this document may be reproduced without the prior written permission of Rothamsted Research. Printed: 1988. Rothamsted 1988

87/W/CS/34 Nematicides in Crop Sequence - Potatoes, W. Wheat, S. Barley

Rothamsted Research

Rothamsted Research (1988) *87/W/CS/34 Nematicides in Crop Sequence - Potatoes, W. Wheat, S. Barley* ; Yields Of The Field Experiments 1987, pp 82 - 87 - DOI:

<https://doi.org/10.23637/ERADOC-1-37>

87/W/CS/34

NEMATOCIDES IN CROP SEQUENCE

Object: To study the effects of a range of nematocides on the incidence of *Globodera rostochiensis* and the yield of potatoes. Residual effects of previous treatments are studied in wheat and barley - Woburn Great Hill II and III.

Sponsor: A.G. Whitehead.

The 19th year, potatoes, w. wheat, s. barley.

For previous years see 71/W/CS/34(t), 72/W/CS/34(t) and 73-86/W/CS/34.

Design: 4 series of 3 blocks of 10 plots.

Whole plot dimensions: 4.27 x 9.14.

Treatments: The experiment has four series with the following cropping:-

| | 1970 | 71 | 72 | 73 | 74 | 75 | 76 | 77 | 78 | 79 | 80 | 81 | 82 | 83 | 84 | 85 | 86 | 87 |
|------------|------|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|
| Series I | P | P* | SB | B | P | P* | W | B | P | P* | B | B | P | P* | W | B | P | P |
| Series II | P | P | P* | SB | B | P | P* | W | B | P | P* | W | B | P | P* | B | B | P |
| Series III | B | P | P | P* | SB | B | P | P* | W | B | P | P* | W | B | P | P* | W | B |
| Series IV | B | P | P | P | P* | SB | B | P | P* | W | B | P | P* | W | B | P | P* | W |

P = potatoes, SB = sugar beet, B = s. barley, W = w. wheat

* Treatments applied to potatoes, subsequent crops test residual effects. In 1987 planned new treatments were not applied to Series I and yields were not taken.

Treatments to potatoes (Series II): All combinations of:-

1. NEMACIDE[84] Residues of nematocides applied 1984:

ALDICARB
SDS38697
SDS46995

2. RATE Rates of nematocide (a.i.):

SINGLE Single (2.8 kg of aldicarb, 1.4 kg of SDS materials)
DOUBLE Double (5.6 kg of aldicarb, 2.8 kg of SDS materials)
QUAD Quadruple (11.2 kg of aldicarb, 5.6 kg of SDS materials)

plus one untreated plot per block

RATE

NONE

87/W/CS/34

Treatments to s. barley (Series III): All combinations of:-

1. NEMACIDE[85] Residues of nematicides applied 1985:

| | |
|----------|-------------|
| OXAMYL | Oxamyl |
| SDS46995 | 'SDS 46995' |
| THIODICA | Thiodicarb |

2. RATE Rates of nematicide (a.i.):

| | |
|--------|---|
| SINGLE | Single (1.4 kg of 'SDS 46995', 2.8 kg of other materials) |
| DOUBLE | Double (2.8 kg of 'SDS 46995', 5.6 kg of other materials) |
| QUAD | Quadruple (5.6 kg of 'SDS 46995', 11.2 kg of other materials) |

plus one untreated plot per block

RATE

NONE

Treatments to w. wheat (Series IV):

1. NEMACIDE[86] Residues of nematicides and rates (a.i.) applied 1986:

| | |
|----------|--|
| AL 3.3 | Aldicarb at 3.3 kg |
| AL 6.6 | Aldicarb at 6.6 kg |
| AL S 3.3 | Aldicarb, slow release formulation at 3.3 kg |
| AL S 6.6 | Aldicarb, slow release formulation at 6.6 kg |
| ETH 7.5 | Ethoprophos at 7.5 kg |
| MB 5.0 | 'MB 41380' at 5.0 kg |
| MB 7.5 | 'MB 41380' at 7.5 kg |
| MB 10.0 | 'MB 41380' at 10.0 kg |
| OX 5.0 | Oxamyl at 5.0 kg |
| NONE | None |

Standard applications:

- Potatoes (Series I and II): Manures: (10:10:15+4.5 Mg) at 2300 kg. Weedkillers: Glyphosate at 1.4 kg in 200 l to Series II only. Linuron at 1.6 kg in 200 l. Fungicides: Mancozeb at 1.4 kg on four occasions in 200 l, applied with the pirimicarb on the second. Fentin hydroxide at 0.28 kg on two occasions in 200 l. Nematicide: Oxamyl at 5.6 kg to Series I only. Insecticide: Pirimicarb at 0.14 kg. Desiccant: Diquat at 0.80 kg ion in 200 l.
- W. wheat (Series IV): Manures: N at 210 kg as 'Nitram'. Weedkillers: Clopyralid at 0.05 kg, bromoxynil at 0.24 kg with mecoprop at 2.5 kg applied with the tridemorph in 200 l. Fungicides: Tridemorph at 0.52 kg. Triadimenol at 0.062 kg with tridemorph at 0.38 kg in 200 l.
- S. barley (Series III): Manures: (20:10:10) at 630 kg. Weedkillers: Glyphosate at 1.4 kg in 200 l. Clopyralid at 0.05 kg, bromoxynil at 0.24 kg with mecoprop at 2.5 kg applied with the tridemorph in 200 l. Fungicides: Tridemorph at 0.52 kg. Triadimenol at 0.062 kg with tridemorph at 0.38 kg in 200 l.

87/W/CS/34

Seed: Potatoes: Pentland Crown.
W. wheat: Avalon, sown at 210 kg.
S. barley: Klaxon, sown at 160 kg.

Cultivations, etc.:-

- Potatoes (Series I and II): Glyphosate applied (Series II only): 19 Sept, 1986. Ploughed (Series II only): 21 Nov. Spring-tine cultivated: 18 Feb, 1987. NPK Mg applied: 21 Apr. Oxamyl applied (Series I only), rotary cultivated, potatoes planted: 27 Apr. Rotary ridged: 15 May. Linuron applied: 25 May. Mancozeb applied: 24 June, 8 July, 26 July, 5 Aug. Pirimicarb applied: 8 July. Fentin hydroxide applied: 18 Aug, 4 Sept. Desiccant applied: 18 Sept. Haulm mechanically destroyed: 1 Oct. Lifted; (Series I): 12 Oct, (Series II): 13 Oct.
- W. wheat (Series IV): Spring-tine cultivated, rotary harrowed with crumbler attached, seed sown, harrowed: 4 Dec, 1986. N applied: 5 May, 1987. Weedkillers and tridemorph applied: 29 May. Triadimenol and tridemorph applied: 4 July. Combine harvested: 15 Sept.
- S. barley (Series III): Glyphosate applied: 19 Sept, 1986. Ploughed: 21 Nov. Spring-tine cultivated: 18 Feb, 1987. NPK applied, spike harrowed with crumbler attached, seed sown: 19 Mar. Clopyralid, bromoxynil, mecoprop and tridemorph applied: 29 May. Triadimenol and tridemorph applied: 4 July. Combine harvested: 21 Aug.

87/W/CS/34

POTATOES SERIES II

TOTAL TUBERS TONNES/HECTARE

***** Tables of means *****

| | RATE | SINGLE | DOUBLE | QUAD | Mean |
|--------------|------|--------|--------|------|------|
| NEMACIDE[84] | | | | | |
| ALDICARB | | 38.4 | 37.7 | 40.7 | 38.9 |
| SDS38697 | | 29.1 | 38.2 | 38.8 | 35.3 |
| SDS46995 | | 36.1 | 35.2 | 34.8 | 35.4 |
| Mean | | 34.5 | 37.0 | 38.1 | 36.5 |
| RATE NONE | | 30.6 | | | |
| Grand mean | | 35.9 | | | |

*** Standard errors of differences of means ***

| Table | NEMACIDE[84] | RATE NEMACIDE[84] RATE & RATE NONE |
|--------|--------------|--|
| s.e.d. | 1.22 | 1.22 2.11 |

***** Stratum standard errors and coefficients of variation *****

| Stratum | d.f. | s.e. | cv% |
|----------|------|------|-----|
| BLOCK.WP | 18 | 2.58 | 7.2 |

PERCENTAGE WARE 3.81 CM (1.5 INCH) RIDDLE

***** Tables of means *****

| | RATE | SINGLE | DOUBLE | QUAD | Mean |
|--------------|------|--------|--------|------|------|
| NEMACIDE[84] | | | | | |
| ALDICARB | | 86.3 | 83.8 | 86.0 | 85.4 |
| SDS38697 | | 85.9 | 85.1 | 84.7 | 85.2 |
| SDS46995 | | 86.6 | 86.0 | 83.8 | 85.5 |
| Mean | | 86.2 | 84.9 | 84.8 | 85.3 |
| RATE NONE | | 83.7 | | | |
| Grand mean | | 85.2 | | | |

PLOT AREA HARVESTED 0.00130

87/W/CS/34

W. WHEAT SERIES IV

GRAIN TONNES/HECTARE

***** Tables of means *****

| NEMACIDE[86] | |
|--------------|------|
| AL 3.3 | 4.39 |
| AL 6.6 | 4.67 |
| AL S 3.3 | 4.19 |
| AL S 6.6 | 5.36 |
| ETH 7.5 | 4.45 |
| MB 5.0 | 4.39 |
| MB 7.5 | 4.94 |
| MB 10.0 | 5.39 |
| OX 5.0 | 4.60 |
| NONE | 4.21 |
| Mean | 4.66 |

*** Standard errors of differences of means ***

| Table | NEMACIDE[86] |
|--------|--------------|
| s.e.d. | 0.498 |

***** Stratum standard errors and coefficients of variation *****

| Stratum | d.f. | s.e. | cv% |
|---------------------|---------|-------|------|
| BLOCK.WP | 18 | 0.610 | 13.1 |
| GRAIN MEAN DM% | 81.3 | | |
| PLOT AREA HARVESTED | 0.00251 | | |

87/W/CS/34

S. BARLEY SERIES III

GRAIN TONNES/HECTARE

***** Tables of means *****

| | RATE | SINGLE | DOUBLE | QUAD | Mean |
|--------------|------|--------|--------|------|------|
| NEMACIDE[85] | | | | | |
| OXAMYL | | 5.10 | 4.79 | 4.86 | 4.92 |
| SDS46995 | | 5.16 | 4.96 | 4.88 | 5.00 |
| THIODICA | | 4.98 | 4.92 | 5.23 | 5.04 |
| Mean | | 5.08 | 4.89 | 4.99 | 4.99 |
| RATE NONE | 4.22 | | | | |
| Grand mean | 4.91 | | | | |

*** Standard errors of differences of means ***

| Table | NEMACIDE[85] | RATE NEMACIDE[85] | RATE & RATE NONE |
|--------|--------------|-------------------|---------------------|
| s.e.d. | 0.237 | 0.237 | 0.411 |

***** Stratum standard errors and coefficients of variation *****

| Stratum | d.f. | s.e. | cv% |
|---------------------|---------|-------|------|
| BLOCK.WP | 18 | 0.504 | 10.3 |
| GRAIN MEAN DM% | 88.1 | | |
| PLOT AREA HARVESTED | 0.00251 | | |