

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

ARC, Institute of Arable Crops Research
Rothamsted Experimental Station
Harpenden
Herts
SG8 5LR
ENGLAND
UK
The copyright in this document is owned by the Rothamsted Research Ltd
and is a part of the Rothamsted Research Archives. It is made available
here for the purpose of providing a permanent record of the work and
to ensure that the work is accessible to all who wish to use it.
Printed: Rothamsted, Rothamsted 2008

87/R/CS/133 Control of Pathogens - Maize

Rothamsted Research

Rothamsted Research (1988) *87/R/CS/133 Control of Pathogens - Maize* ; Yields Of The Field Experiments 1987, pp 88 - 89 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/CS/133

CONTROL OF PATHOGENS

Object: To study the effects of a range of chemicals on the incidence of pathogens and yield of continuous maize - Long Hoos VI/VII 6.

Sponsors: A.J. Barnard, D.J. Hooper, D. Hornby, R.T. Plumb.

The 14th year, forage maize.

For previous years see 74-86/R/CS/133.

Design: 3 randomised blocks of 9 plots.

Whole plot dimensions: 2.13 x 18.3.

Treatments:-

CHEM RES	Residual effect of chemicals applied annually in previous years (except as stated), none in 1987:
NONE	None (2 plots per block)
ALDICARB	Aldicarb, 4.5 kg as granules to seedbed
BENOMYL	Benomyl, 11.2 kg as dust to seedbed
DAZOMET	Dazomet, 450 kg as granules in early spring (not applied 1975, 1979, 1981 and 1986)
PERMETH	Permethrin, as foliar spray (in 1979, 1984 and 1985 only)
PHORATE	Phorate, 1.68 kg as granules to seedbed
PIRIMICA	Pirimicarb, as foliar spray (in 1979, 1984 and 1985 only)
BE+DA+PH	Benomyl + dazomet (not applied 1975, 1979, 1981, and 1986) + phorate, at above rates and times

Basal applications: Manure: 'Nitro-Chalk' at 550 kg. Weedkiller: Atrazine at 1.7 kg in 220 l.

Seed: Fronica, sown at 100,000 seeds per hectare.

Cultivations, etc.: - Ploughed: 24 Dec, 1986. N applied, spring-tine cultivated twice, seed sown: 6 May, 1987. Rolled, weedkiller applied: 7 May. Harvested by hand: 3 Nov.

87/R/CS/133

FORAGE DRY MATTER TONNES/HECTARE

***** Tables of means *****

CHEM RES	
NONE	9.04
ALDICARB	9.30
BENOMYL	8.98
DAZOMET	10.06
PERMETH	9.05
PHORATE	9.00
PIRIMICA	8.85
BE+DA+PH	8.61
Mean	9.10

*** Standard errors of differences of means ***

Table	CHEM RES
s.e.d.	0.763 min.rep
	0.661 max-min

	CHEM RES
max-min	NONE v any of remainder
min.rep	any of remainder

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	17	0.935	10.3
FORAGE MEAN DM%	20.8		
PLOT AREA HARVESTED	0.00065		