

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1987

[Full Table of Content](#)

ARC, Institute of Arable Crops Research
Rothamsted Experimental Station
Harpenden
Herts
AL5 2JQ
UK
The copyright in this document is held by the Rothamsted Research and is licensed to the National Institute for Research in Food Safety and Food Quality (NIFFSQ) for the purpose of making this document available on the ERADOC platform. All other rights reserved.
Printed: 2018-06-01
Rothamsted 2018

87/R/CS/212 Seasonal Effects of Take-all - W. Wheat, S. Beans

Rothamsted Research

Rothamsted Research (1988) *87/R/CS/212 Seasonal Effects of Take-all - W. Wheat, S. Beans* ; Yields Of The Field Experiments 1987, pp 95 - 96 - DOI: <https://doi.org/10.23637/ERADOC-1-37>

87/R/CS/212

SEASONAL EFFECTS OF TAKE-ALL

Object: To study the incidence of take-all (*Gaeumannomyces graminis*) in continuous w. wheat and in first, second and third w. wheats after a break - Great Harpenden I.

Sponsors: D. Hornby, R.J. Gutteridge.

The tenth year, s. beans, w. wheat.

For previous years see 78-86/R/CS/212.

Design: 3 randomised blocks of 8 plots.

Whole plot dimensions: 5.33 x 10.7.

Treatments:

PREVCROP Previous crops before w. wheat 1987:

	1984	1985	1986
CONT W	W	W	W
FIRST W	W	BE	W
SECOND W	BE	W	W
THIRD W	W	W	W
BEANS 1	W	W	BE (duplicated)
BEANS 2	W	BE	BE

BE = s. beans, W = w. wheat

NOTE: One additional crop sequence was in s. beans 1987, yields not taken.

Standard applications:

W. wheat: Manures: Chalk at 5.0 t. 'Nitram' at 410 kg. Weedkillers: Isoproturon at 2.5 kg in 200 l. Fluroxypyr at 0.20 kg with clopyralid at 0.05 kg and bromoxynil at 0.24 kg in 500 l.

Seed: W. wheat: Avalon, sown at 180 kg.
S. beans: Minden, sown at 260 kg.

Cultivations, etc.:-

Both crops: Chalk applied: 24 Sept, 1986. Ploughed: 29 Sept. Rotary harrowed: 30 Sept.

W. wheat: Seed sown: 1 Oct, 1986. Isoproturon applied: 14 Apr, 1987. N applied: 17 Apr. Remaining weedkillers applied: 21 Apr. Combine harvested: 1 Sept.

S. beans: Rotary harrowed: 20 Mar, 1987. Seed sown: 31 Mar. Hand hoed: 27 May. Combine harvested: 21 Sept.

NOTE: Take-all was assessed in soil and in w. wheat roots.

87/R/CS/212

GRAIN TONNES/HECTARE

***** Tables of means *****

PREVCROP	CONT W	FIRST W	SECOND W	THIRD W	BEANS 1	BEANS 2	Mean
	4.63	4.90	4.68	5.07	5.24	6.00	5.11

*** Standard errors of differences of means ***

Table	PREVCROP
s.e.d.	0.351 min.rep
	0.304 max-min

	PREVCROP
max-min	BEANS 1 v any of the remainder
min.rep	any of the remainder

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	13	0.430	8.4

GRAIN MEAN DM% 85.4

PLOT AREA HARVESTED 0.00303