


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1988

[Full Table of Content](#)



88/R/BE/1 Control of Chocolate Spot and Rust - W. Beans

Rothamsted Research

Rothamsted Research (1989) *88/R/BE/1 Control of Chocolate Spot and Rust - W. Beans* ; Yields Of The Field Experiments 1988, pp 198 - 199 - DOI: <https://doi.org/10.23637/ERADOC-1-43>

88/R/BE/1

WINTER BEANS

CONTROL OF CHOCOLATE SPOT AND RUST

Object: To compare maneb plus mancozeb with benomyl plus chlorothalonil for the control of chocolate spot (*Botrytis* spp.) and rust (*Uromyces viciae-fabae*) on w. beans sown at two densities - New Zealand.

Sponsors: J. McEwen, D.P. Yeoman.

Design: 2 randomised blocks of 18 plots.

Whole plot dimensions: 6.0 x 10.0.

Treatments: All combinations of:-

1. **SEEDRATE** Seeds sown per square metre:

 12
 36

2. **CS FUNG** Fungicides applied to control chocolate spot until first rust pustules seen:

 NONE None
 BEN+CHL Benomyl at 0.50 kg plus chlorothalonil at 1.0 kg with a wetting agent ('Enhance' at 0.03 l) on 20 June, 1988
 MAN+MANC Maneb plus mancozeb each at 0.80 kg on 20 June

3. **RUSTFUNG** Fungicides applied to control rust, first applied as soon as rust pustules seen:

 NONE None
 BEN+CHL Benomyl at 0.50 kg plus chlorothalonil at 1.0 kg with a wetting agent ('Agral' at 0.06 l) on 29 June
 MAN+MANC Maneb plus mancozeb each at 0.80 kg on 29 June

NOTE: All spray treatments were applied in 200 l.

Basal applications: Weedkillers: Paraquat at 0.60 kg ion in 200 l.
Propyzamide at 0.85 kg with simazine at 1.2 kg in 200 l.
Insecticide: Deltamethrin at 0.0079 kg in 200 l on two occasions.

Seed: Bourdon, dressed with thiram.

Cultivations, etc.:- Paraquat applied: 22 Oct, 1987. Seed broadcast and ploughed in: 23 Oct. Disced: 26 Oct. Remaining weedkillers applied: 5 Nov. Insecticide applied: 26 Apr, 1988, 17 May. Combine harvested: 3 Oct. Previous crops: W. wheat 1986 and 1987.

NOTE: Establishment counts were made in autumn, disease assessments were made in July and August and components of yield were measured at maturity.

88/R/BE/1

GRAIN TONNES/HECTARE

***** Tables of means *****

CS FUNG SEEDRATE	NONE	BEN+CHL	MAN+MANC	Mean
12	5.43	6.68	5.89	6.00
36	5.70	6.14	6.21	6.01
Mean	5.56	6.41	6.05	6.01

RUSTFUNG SEEDRATE	NONE	BEN+CHL	MAN+MANC	Mean
12	5.82	5.72	6.47	6.00
36	5.77	6.20	6.06	6.01
Mean	5.80	5.96	6.27	6.01

RUSTFUNG CS FUNG	NONE	BEN+CHL	MAN+MANC	Mean
NONE	5.12	5.27	6.30	5.56
BEN+CHL	6.43	6.55	6.24	6.41
MAN+MANC	5.84	6.06	6.25	6.05
Mean	5.80	5.96	6.27	6.01

SEEDRATE	RUSTFUNG CS FUNG	NONE	BEN+CHL	MAN+MANC
12	NONE	4.89	4.89	6.51
	BEN+CHL	7.11	6.64	6.28
	MAN+MANC	5.45	5.63	6.61
36	NONE	5.34	5.65	6.10
	BEN+CHL	5.75	6.47	6.20
	MAN+MANC	6.23	6.48	5.90

*** Standard errors of differences of means ***

SEEDRATE	CS FUNG	RUSTFUNG	SEEDRATE CS FUNG
0.160	0.196	0.196	0.277
SEEDRATE RUSTFUNG	CS FUNG RUSTFUNG	SEEDRATE CS FUNG RUSTFUNG	
0.277	0.339	0.480	

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	17	0.480	8.0
GRAIN MEAN DM%	77.5		
PLOT AREA HARVESTED	0.00330		