

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1988

[Full Table of Content](#)



88/R/BE/2 Varieties - W. Beans

Rothamsted Research

Rothamsted Research (1989) *88/R/BE/2 Varieties - W. Beans* ; Yields Of The Field Experiments 1988, pp 200 - 201 - DOI: <https://doi.org/10.23637/ERADOC-1-43>

88/R/BE/2

WINTER BEANS

VARIETIES

Object: To compare three varieties of winter beans - New Zealand.

Sponsors: J. McEwen, D.P. Yeoman.

Design: 6 randomised blocks of 3 plots.

Whole plot dimensions: 6.0 x 10.0.

Treatments:

VARIETY Varieties, sown at 24 seeds per square metre on 23 Oct, 1987:

BANNER
BOURDON
PUNCH

Basal applications: Weedkillers: Paraquat at 0.60 kg ion in 200 l. Propyzamide at 0.85 kg with simazine at 1.2 kg in 200 l. Fungicides: Benomyl at 0.50 kg with chlorothalonil at 1.0 kg and a wetting agent ('Enhance' at 0.03 l) in 200 l. Insecticide: Deltamethrin at 0.0079 kg in 200 l on two occasions.

Cultivations, etc.:- Paraquat applied: 22 Oct, 1987. Seed broadcast and ploughed in: 23 Oct. Disced: 26 Oct. Remaining weedkillers applied: 5 Nov. Insecticide applied: 26 Apr, 1988, 17 May. Fungicides with wetting agent applied: 20 June. Combine harvested: 3 Oct. Previous crops: W. wheat 1986 and 1987.

NOTE: Establishment counts were made in autumn and components of yield measured at maturity.

88/R/BE/2

GRAIN TONNES/HECTARE

***** Tables of means *****

VARIETY	BANNER	BOURDON	PUNCH	Mean
	5.24	5.47	5.98	5.57

*** Standard errors of differences of means ***

VARIETY
0.182

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
---------	------	------	-----

BLOCK.WP	10	0.315	5.7
----------	----	-------	-----

GRAIN MEAN DM% 79.8

PLOT AREA HARVESTED 0.00330