


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1988

[Full Table of Content](#)



---

### **88/R/BE/4 Varieties, Row Spacing and Plant Health - S. Beans**

#### **Rothamsted Research**

Rothamsted Research (1989) *88/R/BE/4 Varieties, Row Spacing and Plant Health - S. Beans* ; Yields Of The Field Experiments 1988, pp 202 - 203 - DOI: <https://doi.org/10.23637/ERADOC-1-43>

88/R/BE/4

SPRING BEANS

VARIETIES, ROW SPACING AND PLANT HEALTH

**Object:** To compare four varieties of spring-sown field beans at two row spacings and two amounts of pest and disease control - Long Hoos I/II.

**Sponsors:** J. McEwen, D.P. Yeoman.

**Design:** 3 randomised blocks of 16 plots.

**Whole plot dimensions:** 6.0 x 10.0.

**Treatments:** All combinations of:-

1. **VARIETY** Varieties sown at 60 seeds per square metre:  
  
ALFRED  
MINDEN  
TICOL  
TROY
2. **ROW SPAC** Spacing between rows (cm):  
  
12  
48
3. **PATHCONT** Pest and pathogen control:  
  
STANDARD None  
ENHANCED Economic enhanced to include:  
Deltamethrin at 0.0079 kg in 200 l on 26 Apr, 1988 and 17 May.  
Pirimicarb at 0.14 kg in 200 l on 20 May.  
Benomyl at 0.50 kg with chlorothalonil at 1.0 kg and a wetting agent ('Enhance' at 0.03 l) in 200 l on 20 June.  
Maneb at 0.80 kg and mancozeb at 0.80 kg in 200 l on 29 June.

**Basal applications:** Weedkillers: Trietazine at 1.2 kg and simazine at 0.17 kg in 200 l.

**Cultivations, etc.:-** Ploughed: 1 Dec, 1987. Heavy spring-tine cultivated twice: 7 Mar, 1988. Rotary harrowed, seed sown: 9 Mar. Weedkillers applied: 17 Mar. Combine harvested: 20 Sept. Previous crops: W. wheat 1986 and 1987.

**NOTE:** Establishment counts were made. Disease assessments were made in July and August. At maturity, plant heights, lodging and components of yield were assessed.

88/R/BE/4

GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

ROW SPAC	12	48	Mean
<b>VARIETY</b>			
ALFRED	6.17	5.99	6.08
MINDEN	5.84	5.56	5.70
TICOL	4.54	4.68	4.61
TROY	4.86	5.28	5.07
Mean	5.35	5.38	5.36

PATHCONT	STANDARD	ENHANCED	Mean
<b>VARIETY</b>			
ALFRED	4.90	7.26	6.08
MINDEN	5.30	6.10	5.70
TICOL	3.97	5.25	4.61
TROY	4.33	5.81	5.07
Mean	4.62	6.10	5.36

PATHCONT	STANDARD	ENHANCED	Mean
<b>ROW SPAC</b>			
12	4.70	6.00	5.35
48	4.54	6.21	5.38
Mean	4.62	6.10	5.36

	ROW SPAC	12	48		ROW SPAC	12	48
<b>VARIETY</b>	<b>PATHCONT</b>	STANDARD	ENHANCED	STANDARD	ENHANCED		
ALFRED		5.16	7.18	4.64	7.34		
MINDEN		5.45	6.22	5.14	5.98		
TICOL		3.88	5.21	4.07	5.28		
TROY		4.33	5.39	4.33	6.22		

\*\*\* Standard errors of differences of means \*\*\*

VARIETY	ROW SPAC	PATHCONT	VARIETY
			ROW SPAC
0.171	0.121	0.121	0.242

VARIETY	ROW SPAC	VARIETY
PATHCONT	PATHCONT	ROW SPAC
		PATHCONT
0.242	0.171	0.342

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	30	0.419	7.8
GRAIN MEAN DM%	84.3		
PLOT AREA HARVESTED	ROW SPAC 12	0.00330	
	ROW SPAC 48	0.00312	