


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1988

[Full Table of Content](#)



### 88/R/RA/2 Seed Rates and Row Spacings - W. Oilseed Rape

#### Rothamsted Research

Rothamsted Research (1989) *88/R/RA/2 Seed Rates and Row Spacings - W. Oilseed Rape* ; Yields Of The Field Experiments 1988, pp 218 - 219 - DOI: <https://doi.org/10.23637/ERADOC-1-43>

88/R/RA/2

WINTER OILSEED RAPE

SEED RATES AND ROW SPACINGS

**Object:** To compare cv. Ariana on a range of row-widths and seed rates - Whittlocks.

**Sponsor:** D.P. Yeoman.

**Design:** 3 randomised blocks of 11 plots.

**Whole plot dimensions:** 3.0 x 15.0.

**Treatments:** All combinations of:-

1. **SEEDRATE**            Seed rates:

4 KG  
6 KG  
8 KG

2. **ROWSPACE**            Row spacings:

17.5 CM  
35 CM  
52.5 CM

plus two extra treatments, sown at 2 kg seed rate:-

**EXTRA**                    Row spacings:

2 KG 35                    35 cm  
2 KG 52.5                   52.5 cm

**Basal applications:** Manures: 'Nitram' at 140 kg and later at 720 kg. Weedkillers: Paraquat at 0.40 kg ion in 200 l. Metazachlor at 1.2 kg in 200 l. Insecticides: Azinphos methyl at 0.40 kg and demeton-s-methyl sulphone at 0.12 kg in 300 l. Desiccant: Diquat at 0.60 kg ion with a wetting agent ('Enhance' at 0.50 l) in 520 l.

**Seed:** Ariana, dressed gamma HCH, thiram and fenpropimorph.

**Cultivations, etc.:-** Paraquat applied: 8 Sept, 1987. First N applied, heavy spring-tine cultivated: 14 Sept. Rotary harrowed, seed sown: 15 Sept. Metazachlor applied: 1 Oct. Second N applied: 19 Feb, 1988. Insecticides applied: 18 Apr. Desiccant with wetting agent applied: 20 July. Combine harvested: 27 July. Previous crops: W. wheat 1986, w. barley 1987.

**NOTE:** Plant counts were made at establishment and in spring.

88/R/RA/2

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

ROWSPACE	17.5 CM	35 CM	52.5 CM	Mean
<b>SEEDRATE</b>				
4 KG	2.92	3.83	3.32	3.36
6 KG	3.64	3.16	3.52	3.44
8 KG	3.29	3.00	3.27	3.18
Mean	3.28	3.33	3.37	3.33

EXTRA	2 KG	35	2 KG	52.5	Mean
	3.45		3.97		3.71

GRAND MEAN 3.40

\*\*\* Standard errors of differences of means \*\*\*

SEEDRATE	ROWSPACE	SEEDRATE ROWSPACE & EXTRA
0.206	0.206	0.356

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	20	0.436	12.8
MEAN DM%	72.2		
PLOT AREA HARVESTED	0.00345		