


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1988

[Full Table of Content](#)



### 88/R/P/1 and 88/W/P/1 Varieties - Potatoes

#### Rothamsted Research

Rothamsted Research (1989) *88/R/P/1 and 88/W/P/1 Varieties - Potatoes* ; Yields Of The Field Experiments 1988, pp 250 - 252 - DOI: <https://doi.org/10.23637/ERADOC-1-43>

88/R/P/1 and 88/W/P/1

POTATOES

VARIETIES

**Object:** To compare the quality and yield of some of the newer varieties of potato with current standards on two soil types - Rothamsted Pastures (R), Woburn White Horse (W).

**Sponsor:** R. Moffitt.

**Design:** 4 randomised blocks of 8 plots.

**Whole plot dimensions:** 3.0 x 7.0.

**Treatments:**

VARIETY	Varieties:
CARA	Cara
CROWN	Pentland Crown
DESIREE	Desiree
DE+CAO	Desiree given calcium peroxide in the ridge at planting
ESTIMA	Estima
KINGSTON	Kingston
ROMANO	Romano
WILJA	Wilja

**NOTE:** Calcium peroxide was applied as granules at a rate of 370 kg (10 g per tuber) applied to planted tuber before covering.

**Basal applications:**

Pastures (R): Manures: (0:18:36) at 690 kg. FYM at 35 t. (10:10:15+4.5 Mg) at 1960 kg. Weedkiller: Linuron at 1.65 kg in 260 l. Fungicides: Dithiocarbamate complex at 1.4 kg in 200 l applied twice. Mancozeb at 1.4 kg in 200 l on five occasions applied with the insecticide on the first two. Fentin hydroxide at 0.27 kg in 200 l. Insecticide: Pirimicarb at 0.14 kg. Desiccant: Diquat at 0.80 kg ion in 200 l.

White Horse (W): Manures: (0:18:36) at 690 kg. (10:10:15+4.5 Mg) at 2400 kg. Weedkillers: Paraquat at 0.40 kg ion in 200 l. Linuron at 1.5 kg with paraquat at 0.40 kg ion in 220 l. Fungicides: Mancozeb at 1.4 kg in 220 l on five occasions applied with the insecticide on the first, second and fifth occasions. Fentin hydroxide at 0.28 kg in 220 l. Nematicide: Oxamyl at 5.5 kg. Insecticide: Pirimicarb at 0.14 kg. Desiccant: Diquat at 0.80 kg ion in 400 l.

**Cultivations, etc.:-**

Pastures (R): PK applied: 13 Oct, 1987. FYM applied: 10 Dec. Ploughed: 17 Dec. NPK Mg applied: 6 Apr, 1988. Rotary harrowed, ridged, hand planted and split back: 11 Apr. Ridged: 26 Apr. Linuron applied: 5 May. Mancozeb and pirimicarb applied: 15 and 30 June. Dithiocarbamate complex applied: 6 and 8 July. Mancozeb applied: 18 July, 1 and 15 Aug. Fentin hydroxide applied: 30 Aug. Haulm mechanically destroyed: 14 Sept. Desiccant applied: 20 Sept. Lifted: 31 Oct. Previous crops: W. wheat 1986, s. beans 1987.

88/R/P/1 and 88/W/P/1

Cultivations, etc.:-

White Horse (W): Paraquat applied: 21 Sept, 1987. PK applied: 25 Sept. NPK Mg applied: 7 Apr, 1988. Oxamyl applied, rotary cultivated and ridged, hand planted, split back ridges: 15 Apr. Rotary ridged, linuron and paraquat applied: 12 May. Mancozeb applied: 15 July, 1 Aug. Mancozeb and pirimicarb applied: 14 June, 7 July and 15 Aug. Fentin hydroxide applied: 30 Aug. Desiccant applied: 15 Sept. Haulm mechanically destroyed: 29 Sept. Lifted: 14 Oct. Previous crops: Mixed w. cereals 1986, w. oats 1987.

NOTE: On Pastures (R) one of the plots of CROWN was severely infested with thistles. An estimated value was used in the analysis.

88/R/P/1 PASTURES (R)

TOTAL TUBERS TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

VARIETY	
CARA	80.7
CROWN	77.7
DESIREE	63.4
DE+CAO	63.5
ESTIMA	74.7
KINGSTON	68.1
ROMANO	66.3
WILJA	67.9
Mean	70.3

\*\*\* Standard errors of differences of means \*\*\*

VARIETY
3.54

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	20	5.00	7.1

PERCENTAGE WARE 3.81 CM (1.5 INCH) RIDDLE

\*\*\*\*\* Tables of means \*\*\*\*\*

VARIETY	
CARA	98.0
CROWN	98.5
DESIREE	97.0
DE+CAO	96.8
ESTIMA	98.3
KINGSTON	97.4
ROMANO	98.0
WILJA	96.8
Mean	97.6

PLOT AREA HARVESTED 0.00105

88/W/P/1 WHITE HORSE (W)

TOTAL TUBERS TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

VARIETY	
CARA	83.3
CROWN	76.3
DESIREE	60.4
DE+CAO	53.6
ESTIMA	72.5
KINGSTON	72.9
ROMANO	62.7
WILJA	74.4
Mean	69.5

\*\*\* Standard errors of differences of means \*\*\*

VARIETY
3.24

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	21	4.59	6.6

PERCENTAGE WARE 3.81 CM (1.5 INCH) RIDDLE

\*\*\*\*\* Tables of means \*\*\*\*\*

VARIETY	
CARA	96.3
CROWN	96.2
DESIREE	94.5
DE+CAO	93.8
ESTIMA	95.2
KINGSTON	95.5
ROMANO	95.3
WILJA	93.7
Mean	95.0

PLOT AREA HARVESTED 0.00105