Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1988



Full Table of Content

88/R/M/1 and 88/W/M/1 Inputs for Winter Cereals - W. Triticale, Wheat, Barley, Rye - Mixed Crops

Rothamsted Research

Rothamsted Research (1989) 88/R/M/1 and 88/W/M/1 Inputs for Winter Cereals - W. Triticale, Wheat, Barley, Rye - Mixed Crops; Yields Of The Field Experiments 1988, pp 253 - 255 - DOI: https://doi.org/10.23637/ERADOC-1-43

88/R/M/1 and 88/W/M/1

MIXED 1

INPUTS FOR WINTER CEREALS

Object: To compare amounts of disease and the yield of triticale with
 those of w. wheat, w. barley and w. rye on two contrasted sites each
 given contrasted amounts of agrochemicals - Rothamsted Highfield VI (R)
 Woburn Far Field II (W).

Sponsors: R.J. Gutteridge, D. Hornby, R.D. Prew (R), P.R. Scott,
W. Hollins, R.L. Gregory (I.P.S.R., Cambridge).

Design: 3 randomised blocks of 10 plots.

Whole plot dimensions: 3.0 x 10.0 (R), 4.0 x 10.0 (W).

Treatments: All combinations of :-

1. CROP VAR	Crop and variety:	(R)	(W)
B PANDA	W. barley, Panda sown at	200 kg,	200 kg
R DOMINT	W. rye, Dominant sown at	120 kg,	140 kg
- T LASKO	W. triticale, Lasko sown at	180 kg,	180 kg
T STATUS	W. triticale, Status sown at	130 kg,	140 kg
- W AVALON	W. wheat, Avalon sown at	180 kg,	190 kg
2. INPUT	Inputs of agrochemicals, in addit	ion to bas	als:
LARGE	High input, 40 kg N early plus 16 (W).	00 kg N lat	er (R) and
	Prochloraz + carbendazim + trider (W).	norph (Apri	1) (R) and
	Fenpropimorph + chlorothalonil (M	May) (R)	
	Propiconazole + tridemorph (May)	(W)	
	Propiconazole + carbendazim (June	e) (W)	
SMALL	Low input, 120 kg N in early Apri summer insecticides.	il. No fun	gicides or

NOTES: (1) Treatments were applied as follows:

Highfield (R): N treatments: 23 Feb, 1988 and 5 Apr.

Prochloraz at 0.40 kg, carbendazim at 0.15 kg with

tridemorph at 0.52 kg in 220 l: 26 Apr. Fenpropimorph

at 0.75 kg with clorothalonil at 1.0 kg in 220 l: 25 May.

Far Field II (W): N treatments: 3 Mar, 1988 and 13 Apr.

Prochloraz at 0.40 kg, carbendazim at 0.15 kg with

tridemorph at 0.52 kg in 220 l: 26 Apr. Propiconazole

at 0.12 kg and tridemorph at 0.25 kg in 220 l: 23 May.

Propiconazole at 0.12 kg with carbendazim at 0.15 kg:

18 June.

(2) The R DOMINT and the T STATUS plots at Rothamsted failed and were cultivated and fallowed in early May 1988.

88/R/M/1 and 88/W/M/1

Basal applications:

Highfield VI (R): Manures: Chalk at 5.0 t. Weedkillers: Glyphosate at 0.27 kg in 200 l. Diclofop-methyl at 0.95 kg in 200 l. Fluroxypyr at 0.20 kg with metsulfuron-methyl at 0.006 kg in 200 l. Insecticide: Fonofos at 1.4 kg in 200 l. Far Field II (W): Weedkillers: Isoxaben at 0.12 kg in 200 l. Diclofop-methyl at 0.57 kg in 200 l. Insecticide: Fonofos at 1.4 kg in 200 1.

Cultivations, etc.:-

Highfield VI (R): Chalk applied: 16 Sept, 1987. Glyphosate applied: 29 Sept. Ploughed: 2 Oct. Rotary harrowed: 5 Oct. Rotary harrowed, seed sown, harrowed: 6 Oct. Diclofop-methyl applied: 18 Nov. Insecticide applied: 14 Jan, 1988. Fluroxypyr and metsulfuron-methyl applied: 25 Apr. Combine harvested: 4 Aug (barley), 23 Aug (wheat), 26 Aug (triticale). Previous crops: W. wheat 1986 and 1987.

Far Field II (W): Ploughed: 21 Sept, 1987. Rotary cultivated with crumbler attached, seed sown: 1 Oct. Harrowed and rolled: 2 Oct. Isoxaben applied: 23 Oct. Diclofop-methyl applied: 24 Oct. Insecticide applied: 12 Jan, 1988. Combine harvested: 3 Aug (barley) 25 Aug (other cereals). Previous crops: W. wheat 1986 and 1987.

88/R/M/1 W.BARLEY, W.TRITICALE, W.WHEAT

GRAIN TONNES/HECTARE

**** Tables of means ****

INPUT CROP VAR	LARGE	SMALL	Mean
B PANDA	8.40	6.44	7.42
T LASKO	8.27	6.18	7.22
W AVALON	6.88	5.87	6.37
Mean	7.85	6.16	7.01

*** Standard errors of differences of means ***

CROP VAR	INPUT	CROP VAR
		INPUT
0.458	0.374	0.647

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	CV%
BLOCK.WP	10	0.793	11.3

GRAIN MEAN DM% 81.9

PLOT AREA HARVESTED 0.00272

88/W/M/1 W.BARLEY, W.TRITICALE, W.WHEAT

GRAIN TONNES/HECTARE

***** Tables of means *****

INPUT	LARGE	SMALL	Mean
CROP VAR			
B PANDA	6.99	4.52	5.76
R DOMINT	8.15	7.09	7.62
T LASKO	3.90	3.80	3.85
T STATUS	5.22	2.95	4.09
W AVALON	4.26	2.77	3.52
Mean	5.71	4.23	4.97

*** Standard errors of differences of means ***

CROP VAR	INPUT	CROP VAR
		INPUT
0.354	0.224	0.500

**** Stratum standard errors and coefficients of variation ****

Stratum d.f. s.e. cv% BLOCK.WP 18 0.613 12.3

GRAIN MEAN DM% 81.3

PLOT AREA HARVESTED 0.00271