


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1988

[Full Table of Content](#)



### 88/R/M/6 Factors Affecting Eyespot - W. Wheat, Barley - Mixed Crops

#### Rothamsted Research

Rothamsted Research (1989) *88/R/M/6 Factors Affecting Eyespot - W. Wheat, Barley - Mixed Crops* ;  
Yields Of The Field Experiments 1988, pp 259 - 262 - DOI:

<https://doi.org/10.23637/ERADOC-1-43>

88/R/M/6

MIXED 6

**FACTORS AFFECTING EYESPOT**

**Object:** To study eyespot (*Pseudocercospora herpotrichoides*) development after inoculation with different pathotypes in relation to host crop and seed rate - Great Knott II.

**Sponsors:** A. Goulds, B.D.L. Fitt.

**Design:** 2 randomised blocks of 12 plots split into 4.

**Whole plot dimensions:** 3.0 x 37.0.

**Treatments:** All combinations of:-

Whole plots

1. **W CEREAL** Winter cereals sown on 30 October, 1987:  
BARLEY Winter barley cv. Opera  
WHEAT Winter wheat cv. Avalon
2. **SEEDRATE** Seed rates (seeds per square metre):  
NORMAL Normal - 300 barley, 400 wheat  
HALF N Half normal - 150 barley, 200 wheat
3. **INOCULUM** Inoculation with different eyespot pathogen types:  
NONE None  
RYE INOC Rye type  
WHE INOC Wheat type

Sub plots

4. **FUNGTIME** Times of applying prochloraz at 0.40 kg and carbendazim at 0.15 kg in 220 l:  
NONE None  
EARLY Sprayed at growth stage 30/31 on 26 Apr, 1988  
LATE Sprayed at growth stage, 33/37 wheat, 41/49 barley on 20 May

**NOTES:** (1) One additional sub-plot in each whole plot was systematically arranged for sampling, yields not taken.  
(2) Strains of wheat and rye type inoculum were colonised on oat seed and broadcast within two weeks of emergence.

**Basal applications:** Manures: Chalk at 5.0 t. 'Nitram' at 120 kg and later at 250 kg. Weedkillers: Paraquat at 0.60 kg ion in 200 l. Fluroxypyr at 0.20 kg with clopyralid at 0.07 kg and bromoxynil at 0.34 kg with the tridemorph in 200 l. Fungicides: Tridemorph at 0.52 kg. Chlorothalonil at 1.0 kg in 200 l.

88/R/M/6

**Cultivations, etc.:-** Ploughed: 12 Aug, 1987. Discd: 17 Aug. Chalk applied: 16 Sept. Paraquat applied: 28 Oct. Rotary harrowed, seed sown: 30 Oct. First N applied: 2 Mar, 1988. Second N applied 22 Apr. Remaining weedkillers with tridemorph applied: 10 May. Chlorothalonil applied: 6 June. Combine harvested: 4 Aug (barley), 22 Aug (wheat). Previous crops: W. barley 1986, w. oilseed rape 1987.

**NOTE:** Eyespot was assessed on plants at weekly intervals from early March until early August.

**GRAIN TONNES/HECTARE**

\*\*\*\*\* Tables of means \*\*\*\*\*

<b>SEEDRATE</b>	<b>NORMAL</b>	<b>HALF N</b>	<b>Mean</b>	
<b>W CEREAL</b>				
BARLEY	6.93	6.58	6.76	
WHEAT	8.96	8.26	8.61	
Mean	7.94	7.42	7.68	
<b>INOCULUM</b>	<b>NONE</b>	<b>RYE INOC</b>	<b>WHE INOC</b>	<b>Mean</b>
<b>W CEREAL</b>				
BARLEY	6.74	6.81	6.71	6.76
WHEAT	8.62	8.55	8.66	8.61
Mean	7.68	7.68	7.69	7.68
<b>INOCULUM</b>	<b>NONE</b>	<b>RYE INOC</b>	<b>WHE INOC</b>	<b>Mean</b>
<b>SEEDRATE</b>				
NORMAL	7.97	8.04	7.82	7.94
HALF N	7.39	7.32	7.55	7.42
Mean	7.68	7.68	7.69	7.68
<b>FUNGTIME</b>	<b>NONE</b>	<b>EARLY</b>	<b>LATE</b>	<b>Mean</b>
<b>W CEREAL</b>				
BARLEY	6.62	6.82	6.82	6.76
WHEAT	8.42	8.64	8.76	8.61
Mean	7.52	7.73	7.79	7.68
<b>FUNGTIME</b>	<b>NONE</b>	<b>EARLY</b>	<b>LATE</b>	<b>Mean</b>
<b>SEEDRATE</b>				
NORMAL	7.78	8.02	8.04	7.94
HALF N	7.27	7.45	7.55	7.42
Mean	7.52	7.73	7.79	7.68
<b>FUNGTIME</b>	<b>NONE</b>	<b>EARLY</b>	<b>LATE</b>	<b>Mean</b>
<b>INOCULUM</b>				
NONE	7.61	7.73	7.70	7.68
RYE INOC	7.43	7.80	7.82	7.68
WHE INOC	7.53	7.66	7.87	7.69
Mean	7.52	7.73	7.79	7.68

88/R/M/6

GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

		INOCULUM	NONE	RYE INOC	WHE INOC
<b>W CEREAL</b>	<b>SEEDRATE</b>				
BARLEY	NORMAL		6.97	7.03	6.78
	HALF N		6.52	6.59	6.65
WHEAT	NORMAL		8.97	9.05	8.86
	HALF N		8.26	8.05	8.46
		<b>FUNGTIME</b>	<b>NONE</b>	<b>EARLY</b>	<b>LATE</b>
<b>W CEREAL</b>	<b>SEEDRATE</b>				
BARLEY	NORMAL		6.78	7.06	6.93
	HALF N		6.46	6.57	6.72
WHEAT	NORMAL		8.77	8.97	9.15
	HALF N		8.08	8.32	8.38
		<b>FUNGTIME</b>	<b>NONE</b>	<b>EARLY</b>	<b>LATE</b>
<b>W CEREAL</b>	<b>INOCULUM</b>				
BARLEY	NONE		6.64	6.72	6.88
	RYE INOC		6.57	6.98	6.89
	WHE INOC		6.67	6.76	6.71
WHEAT	NONE		8.58	8.75	8.52
	RYE INOC		8.29	8.62	8.75
	WHE INOC		8.39	8.56	9.02
		<b>FUNGTIME</b>	<b>NONE</b>	<b>EARLY</b>	<b>LATE</b>
<b>SEEDRATE</b>	<b>INOCULUM</b>				
NORMAL	NONE		7.75	8.09	8.07
	RYE INOC		7.91	8.06	8.16
	WHE INOC		7.67	7.89	7.89
HALF N	NONE		7.47	7.37	7.32
	RYE INOC		6.95	7.54	7.48
	WHE INOC		7.39	7.43	7.84
		<b>FUNGTIME</b>	<b>NONE</b>	<b>EARLY</b>	<b>LATE</b>
<b>W CEREAL</b>	<b>SEEDRATE</b>	<b>INOCULUM</b>			
BARLEY	NORMAL	NONE	6.76	7.11	7.04
		RYE INOC	6.93	7.21	6.97
		WHE INOC	6.67	6.88	6.79
	HALF N	NONE	6.51	6.32	6.71
		RYE INOC	6.21	6.75	6.81
		WHE INOC	6.67	6.65	6.63
WHEAT	NORMAL	NONE	8.74	9.08	9.11
		RYE INOC	8.90	8.91	9.34
		WHE INOC	8.68	8.91	9.00
	HALF N	NONE	8.43	8.42	7.94
		RYE INOC	7.68	8.33	8.15
		WHE INOC	8.11	8.21	9.05

88/R/M/6

GRAIN TONNES/HECTARE

\*\*\* Standard errors of differences of means \*\*\*

	W CEREAL	SEEDRATE	INOCULUM	FUNGTIME
	0.090	0.090	0.110	0.086

	W CEREAL	W CEREAL	SEEDRATE	W CEREAL
	SEEDRATE	INOCULUM	INOCULUM	FUNGTIME
	0.127	0.156	0.156	0.134

Except when comparing means with the same level(s) of  
W CEREAL 0.122

	SEEDRATE	INOCULUM	W CEREAL	W CEREAL
	FUNGTIME	FUNGTIME	SEEDRATE	SEEDRATE
			INOCULUM	FUNGTIME
	0.134	0.164	0.220	0.190

Except when comparing means with the same level(s) of  
SEEDRATE 0.122  
INOCULUM 0.150  
W CEREAL. SEEDRATE 0.173

	W CEREAL	SEEDRATE	W CEREAL
	INOCULUM	INOCULUM	SEEDRATE
	FUNGTIME	FUNGTIME	INOCULUM
			FUNGTIME
	0.233	0.233	0.329

Except when comparing means with the same level(s) of  
W CEREAL. INOCULUM 0.212  
SEEDRATE. INOCULUM 0.212  
W CEREAL. SEEDRATE. INOCULUM 0.299

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	11	0.220	2.9
BLOCK.WP.SP	24	0.299	3.9

GRAIN MEAN DM% 80.5

SUB PLOT AREA HARVESTED 0.00235