

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1988

[Full Table of Content](#)



88/R/RN/8 Cultivation/WEEDKILLER - W. Barley

Rothamsted Research

Rothamsted Research (1989) *88/R/RN/8 Cultivation/WEEDKILLER - W. Barley* ; Yields Of The Field Experiments 1988, pp 59 - 60 - DOI: <https://doi.org/10.23637/ERADOC-1-43>

88/R/RN/8

CULTIVATION/WEEDKILLER

Object: To study the long-term effects of different methods of primary cultivation on a sequence of crops; weedkillers were also tested until 1981 - Great Harpenden I.

Sponsor: R. Moffitt.

The 28th year, w. barley.

For previous years see 'Details' 1967 and 1973 and 74-87/R/RN/8.

Design: 2 randomised blocks of 12 plots.

Whole plot dimensions: 12.8 x 12.2.

Treatments: All combinations of:-

Whole plots

1. **CLT CHOP** Primary cultivations annually; straw chopped since 1985:

PLOUGH	Ploughed: 18 Aug, 1987
ROTA DIG	Cultivated by rotary digger: 18 Aug
DEEPTINE	Deep-tine cultivated, twice: 18 Aug

2. **SUBSOIL[82]** Subsoiling in September 1982:

NONE	None
CNVNTIAL	Conventional vertical tine
PARAPLOW	'Paraplow'

XTR BURN	plus three extra treatments with straw burnt since 1985, direct drilled until 1984, sprayed with paraquat at 0.60 kg ion in 500 l on 19 Aug, 1987 in addition to basal cultivating, differing in subsoiling in September 1982:
-----------------	--

NONE	None
CNVNTIAL	Conventional vertical tine
PARAPLOW	'Paraplow'

NOTES: (1) Straw was chopped on 8 Aug, 1987 and was burnt and disced on XTR BURN on 21 Aug.
(2) The conventional vertical tine subsoiler had tines 76 cm apart and worked at a depth of about 50 cm.
(3) The 'Paraplow' had rigid tines set at a 45 degree angle. The tip of each tine was in line with the attachment of an adjacent tine. The tines were 51 cm apart and worked at a depth of about 38 cm.

Basal applications: Manures: 'Nitram' at 120 kg followed by 480 kg.
Weedkillers: Glyphosate at 0.27 kg in 200 l. Chlortoluron at 3.5 kg in 200 l.

Seed: Igri, sown at 150 kg.

88/R/RN/8

Cultivations, etc.:- Glyphosate applied: 22 Sept, 1987. Heavy spring-tine cultivated, rotary harrowed, seed sown: 2 Oct. Chlortoluron applied: 6 Nov. First N applied: 1 Mar, 1988. Second N applied: 7 Apr. Combine harvested: 3 Aug.

GRAIN TONNES/HECTARE

***** Tables of means *****

SUBSOIL[82]	NONE	CNVNTIAL	PARAPLOW	Mean
CLT CHOP				
PLOUGH	5.81	5.37	5.52	5.57
ROTA DIG	5.41	4.85	5.58	5.28
DEEPTINE	5.40	5.35	5.12	5.29
Mean	5.54	5.19	5.40	5.38

XTR BURN	NONE	CNVNTIAL	PARAPLOW	Mean
	5.54	5.21	5.62	5.46

Grand mean 5.40

*** Standard errors of differences of means ***

XTR BURN	CLT CHOP	SUBSOIL[82]	CLT CHOP SUBSOIL[82]
0.300	0.173	0.173	0.300

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	11	0.300	5.6

GRAIN MEAN DM% 85.6

PLOT AREA HARVESTED 0.00280