Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1988



Full Table of Content

88/W/RN/13 Intensive Cereals - Potatoes

Rothamsted Research

Rothamsted Research (1989) 88/W/RN/13 Intensive Cereals - Potatoes; Yields Of The Field Experiments 1988, pp 65 - 67 - DOI: https://doi.org/10.23637/ERADOC-1-43

88/W/RN/13

INTENSIVE CEREALS

Object: To study the effects of leys of different duration, following prolonged intensive cereal cropping, on a sequence of arable crops - Woburn Stackyard I.

Sponsors: A.E. Johnston, J. McEwen.

The 23rd year, potatoes.

For previous years see 'Details' 1973 and 74-87/W/RN/13.

Design: 4 randomised blocks of 6 plots split into 6.

Treatments: Until 1977 the experiment tested all phases of the fivecourse rotation: ley, potatoes, cereal, cereal, cereal and continous
cereal. From 1977 to 1980 all phases were cropped with cereal. The
experiment was in two halves, one in which the cereal was w. wheat,
sown on part of the site of the classical continuous wheat experiment
1877-1954 and one in which the cereal was s. barley, sown on part of
the site of the classical continuous barley experiment 1877-1954.
From 1981 the experiment was used to establish grass/clover leys of
different durations for tests on w. wheat in 1987. Plots not in ley
were sown to w. wheat on both halves of the experiment. All leys
were ploughed for 1987 and the site sown to w. wheat. This was
followed in 1988 by potatoes testing all combinations of the
following treatments:

Whole plots

- 1. LEY AGE Length of ley (until ploughing in summer 1986):
 - 1 YEAR
 - 2 YEARS
 - 3 YEARS
 - 4 YEARS
 - 5 YEARS
 - 6 YEARS

Sub plots

- 2. N Nitrogen fertilizer in 1988 (kg N) as 'Nitro-Chalk':
 - 0 70
 - 140
 - 210
 - 280
 - 350

Basal applications: Manures: (0:18:36) at 1400 kg. Mg at 100 kg as kieserite. Weedkillers: Glyphosate at 1.4 kg in 200 l. Linuron at 1.5 kg in 220 l. Fungicides: Mancozeb at 1.4 kg in 220 l on five occasions, applied with the pirimicarb on the first, second and fifth. Fentin hydroxide at 0.28 kg in 220 l. Nematicide: Oxamyl at 5.0 kg. Insecticide: Pirimicarb at 0.14 kg on three occasions. Desiccant: Diquat at 0.80 kg ion in 400 l.

88/W/RN/13

Seed: Pentland Crown.

Cultivations, etc.:- Glyphosate applied: 22 Sept, 1987. Ploughed: 22 Feb, 1988. Heavy spring-tine cultivated: 5 Apr. PK applied: 8 Apr. N treatments applied, oxamyl applied, spring-tine cultivated: 20 Apr. Mg applied, rotary harrowed, potatoes planted: 21 Apr. Rotary ridged, linuron applied: 13 May. Mancozeb applied: 15 July and 1 Aug. Mancozeb applied with pirimicarb: 14 June, 5 July and 15 Aug. Fentin hydroxide applied: 30 Aug. Desiccant applied: 15 Sept. Haulm mechanically destroyed: 29 Sept. Potatoes lifted: 26 Oct.

TOTAL TUBERS TONNES/HECTARE

**** Tables of means ****

N	0	70	140	210	280	350	Mean
LEY AGE							
1 YEAR	37.5	52.6	64.6	60.4	59.4	62.9	56.2
2 YEARS	41.9	57.8	66.2	59.9	64.5	63.3	58.9
3 YEARS	44.9	59.9	64.7	67.6	65.8	63.4	61.0
4 YEARS	46.8	58.1	66.0	67.6	67.9	62.9	61.5
5 YEARS	43.6	61.1	68.6	75.8	69.1	64.9	63.9
6 YEARS	47.5	66.5	64.2	71.1	65.8	67.9	63.8
Mean	43.7	59.3	65.7	67.1	65.4	64.2	60.9

*** Standard errors of differences of means ***

	LEY AGE				N	LEY AGE		
							N	
		2.08		1.	73	4	4.40	
Except when	comparing	means	with	the	same	level	L(s)	of
LEY AGE						4	1.25	

***** Stratum standard errors and coefficients of variation ****

Stratum	d.f.	s.e.	CA &	
BLOCK.WP	15	2.94	4.8	
BLOCK.WP.SP	90	6.01	9.9	

88/W/RN/13

PERCENTAGE WARE 3.81 CM (1.5 INCH) RIDDLE

**** Tables of means ****

N	0	70	140	210	280	350	Mean
LEY AGE							
1 YEAR	94.2	96.5	97.5	96.9	96.9	97.3	96.6
2 YEARS	96.0	97.8	98.0	98.2	98.3	97.5	97.6
3 YEARS	96.7	97.4	98.1	97.3	98.4	97.3	97.5
4 YEARS	95.5	96.9	98.1	97.5	97.5	96.6	97.0
5 YEARS	95.4	97.6	97.9	98.0	98.0	97.5	97.4
6 YEARS	96.1	98.0	97.7	97.7	97.9	97.4	97.5
Mean	95.7	97.4	97.9	97.6	97.8	97.3	97.3

SUB PLOT AREA HARVESTED 0.00090