


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1988

[Full Table of Content](#)



88/R/CS/10 and 88/W/CS/10 Long-term Liming - Linseed

Rothamsted Research

Rothamsted Research (1989) *88/R/CS/10 and 88/W/CS/10 Long-term Liming - Linseed* ; Yields Of The Field Experiments 1988, pp 76 - 80 - DOI: <https://doi.org/10.23637/ERADOC-1-43>

88/R/CS/10 and 88/W/CS/10

LONG TERM LIMING

Object: To study the effects of different amounts of lime and phosphate on the yields and compositions of a sequence of crops. Rothamsted (R) Sawyers I and Woburn (W) Stackyard C.

Sponsor: S.P. McGrath.

The 27th year, linseed.

For previous years see 'Details' 1967, 1973 and 74-87/R&W/CS/10.

Design: 2 randomised blocks of 16 plots split into 2.

Whole plot dimensions: 6.40 x 18.3.

Treatments: All combinations of:-

Whole plots

1. **CHALK** Residual effects of ground chalk (tonnes CaCO₃) (total applied 1962-87):

		Rothamsted total		Woburn total	
R	W	1962-78	1982-87	1962-78	1982-87
0	0	0	0	0	0
15	9	7	8	6	3
24.5	25.5	15	9.5	14	11.5
52.5	45.5	30	22.5	23	22.5

2. **P** Effects of P fertilizer applied:

	Until 1978		1981	1982	1983		1988	
	R	W	R & W	R & W	R	W	R	W
0			0	0	0	0	0	0
P1			0	P1	P1	0	P2	P1
P2			P	P1	0	P2	P2	P1
P3			P	P3	P1	P2	P4	P3

Rates 1981-83 P1, P2, P3, P4 = 25, 50, 75, 100 kg P as superphosphate

Sub plots

3. **MANGNESE** Manganese, cumulative to earlier applications:

0	None
MN	Manganese sprays

- NOTES:** (1) Until 1978 test P was applied cumulatively, rates varied with crop, K was also applied cumulatively, to P1 and P3 plots. Since 1981 K has been applied basally (none in 1986 and 1987).
 (2) On Sawyers I (R) manganese was applied at 0.15 kg Mn, as manganese chelate, in 260 l on 20 June, 1988, repeated in 200 l on 19 July.
 (3) On Stackyard C (W) manganese was applied at 0.16 kg Mn in 220 l on 15 June and at 0.15 kg Mn with sulphur at 0.12 kg S in 220 l on 15 July.

88/R/CS/10 and 88/W/CS/10

Basal applications:

Sawyers I (R): Manures: 'Nitram' at 250 kg. Muriate of potash at 160 kg. Kieserite at 270 kg. Weedkiller: Trifluralin at 1.1 kg in 450 l. Desiccant: Diquat 0.60 kg ion applied with a wetting agent ('Agral' at 0.50 l) in 240 l.

Stackyard C (W): Manures: 'Nitram' at 250 kg. Muriate of potash at 160 kg. Kieserite at 110 kg. Weedkillers: Trifluralin at 0.84 kg in 200 l. Linuron at 0.26 kg in 220 l. Desiccant: Diquat at 0.60 kg ion in 400 l.

Seed: Antares, dressed iprodione and benomyl, sown at 87 kg (R).
Antares, dressed benomyl, sown at 80 kg (W).

Cultivations, etc.:-

Sawyers I (R): P treatments, K and Mg applied: 14 Dec, 1987.
Ploughed: 15 Dec. Heavy spring-tine cultivated: 5 Apr, 1988. N applied: 7 Apr. Weedkiller applied, spring-tine cultivated twice: 12 Apr. Rotary harrowed, seed sown, harrowed: 13 Apr. Desiccant applied: 19 Sept. Combine harvested: 24 Oct.

Stackyard C (W): P treatments, K and Mg applied: 10 Feb, 1988.
Ploughed: 23 Feb. Heavy spring-tine cultivated: 5 Apr. N applied, spike harrowed with crumbler attached, trifluralin applied, spike harrowed with crumbler attached, seed sown: 22 Apr. Linuron applied: 5 May. Desiccant applied: 21 Sept. Combine harvested: 1 Nov.

NOTE: At Rothamsted treatment combinations with **CHALK 0** did not give a measurable yield.

88/R/CS/10 SAWYERS I (R)

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

P	0	P1	P2	P3	Mean
CHALK					
15	2.53	2.57	2.78	2.89	2.69
24.5	2.57	2.90	2.76	2.84	2.77
52.5	2.62	2.81	2.62	2.59	2.66
Mean	2.57	2.76	2.72	2.77	2.70
MANGNESE	0	MN	Mean		
CHALK					
15	2.73	2.65	2.69		
24.5	2.78	2.76	2.77		
52.5	2.65	2.66	2.66		
Mean	2.72	2.69	2.70		

88/R/CS/10 SAWYERS I (R)

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

MANGNESE P	O	MN	Mean
0	2.60	2.55	2.57
P1	2.83	2.69	2.76
P2	2.67	2.77	2.72
P3	2.79	2.76	2.77
Mean	2.72	2.69	2.70

CHALK	MANGNESE P	O	MN
15	0	2.59	2.48
	P1	2.52	2.62
	P2	2.85	2.70
	P3	2.95	2.82
24.5	0	2.55	2.59
	P1	3.07	2.73
	P2	2.62	2.90
	P3	2.88	2.80
52.5	0	2.66	2.57
	P1	2.91	2.71
	P2	2.53	2.72
	P3	2.52	2.65

*** Standard errors of differences of means ***

CHALK	P	MANGNESE	CHALK P
0.136	0.157	0.047	0.272
CHALK MANGNESE	P	CHALK MANGNESE	CHALK MANGNESE
0.148	0.171	0.295	
Except when comparing means with the same level(s) of			
CHALK	0.082		
P		0.094	
CHALK.P			0.163

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	11	0.272	10.1
BLOCK.WP.SP	12	0.163	6.0

GRAIN MEAN DM% 59.8

SUB PLOT AREA HARVESTED 0.00177

88/W/CS/10 STACKYARD C (W)

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

P	0	P1	P2	P3	Mean
CHALK					
0	0.82	1.07	1.72	1.65	1.31
9	2.40	2.90	2.81	2.94	2.76
25.5	2.69	2.69	2.68	3.00	2.77
45.5	2.26	2.48	2.41	2.74	2.47
Mean	2.04	2.29	2.41	2.58	2.33

MANGNESE	0	MN	Mean
CHALK			
0	1.27	1.36	1.31
9	2.77	2.75	2.76
25.5	2.81	2.72	2.77
45.5	2.42	2.53	2.47
Mean	2.32	2.34	2.33

MANGNESE	0	MN	Mean
P			
0	2.00	2.08	2.04
P1	2.34	2.23	2.29
P2	2.36	2.45	2.41
P3	2.56	2.61	2.58
Mean	2.32	2.34	2.33

CHALK	MANGNESE	
	P	MN
0	0	0.74
	P1	0.97
	P2	1.62
	P3	1.73
9	0	2.34
	P1	2.96
	P2	2.81
	P3	2.99
25.5	0	2.82
	P1	2.85
	P2	2.64
	P3	2.93
45.5	0	2.11
	P1	2.59
	P2	2.39
	P3	2.59

88/W/CS/10 STACKYARD C (W)

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

*** Standard errors of differences of means ***

	CHALK	P	MANGNESE	CHALK P
	0.154	0.154	0.057	0.308
	CHALK MANGNESE	P MANGNESE	CHALK P MANGNESE	
	0.174	0.174	0.348	
Except when comparing means with the same level(s) of	CHALK			
	0.114			
P		0.114		
CHALK.P			0.229	

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	15	0.308	13.2
BLOCK.WP.SP	16	0.229	9.8

GRAIN MEAN DM% 83.7

SUB PLOT AREA HARVESTED 0.00185