


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1988

[Full Table of Content](#)



88/R/CS/140 Chemical Reference Plots - S. Barley

Rothamsted Research

Rothamsted Research (1989) *88/R/CS/140 Chemical Reference Plots - S. Barley ; Yields Of The Field Experiments 1988*, pp 85 - 89 - DOI: <https://doi.org/10.23637/ERADOC-1-43>

88/R/CS/140

CHEMICAL REFERENCE PLOTS

Object: To study the persistence in soil of agricultural chemicals applied annually, singly and in combination and their effects on soil microflora and on yield of continuous s. barley - Long Hoos V 3.

Sponsors: R.H. Bromilow, R. Macdonald.

The 15th year, s. barley.

For previous years see 74-87/R/CS/140.

Design: Single replicate of 32 plots.

Whole plot dimensions: 4.06 x 4.57.

Treatments, applied cumulatively except as stated:

All combinations of:-

1. WEEDKLLR Weedkiller in autumn:

NONE	None
GLYPHOS	Glyphosate at 1.4 kg to barley stubble each autumn from 1979 to 1984, at 0.72 kg in 1985, at 0.54 kg in 1986 and at 1.3 kg in 1987

2. FUNGCIDE[1] Fungicide in autumn:

NONE	None
TRIADIM	Triadimefon at 0.25 kg in autumn 1981, 1982, 1984, 1985, 1986 and 1987, 0.28 kg in autumn 1983

3. FUNGCIDE[2] Fungicide in spring:

NONE	None
BENOMYL	Benomyl at 4 kg to seedbed

4. INSCTCDE Insecticide:

NONE	None
CHLORFEN	Chlorfenvinphos at 2 kg to the seedbed

5. NEMACIDE Nematicide:

NONE	None
ALDICARB	Aldicarb at 6 kg to the seedbed

NOTE: Glyphosate and triadimefon were applied in 220 l on 5 Oct, 1987 and 30 Oct respectively. Other treatments were applied on 7 Apr, 1988.

Basal applications: Manure: 'Nitro-Chalk' at 550 kg. Weedkillers: Bentazone at 0.80 kg, dichlorprop at 1.1 kg and MCPA at 0.64 kg in 220 l.

Seed: Doublet, seed not dressed, sown at 160 kg.

88/R/CS/140

Cultivations, etc.:- Ploughed: 27 Nov, 1987. N applied: 29 Feb, 1988. Spring-tine cultivated, seedbed treatments applied, rotary harrowed and seed sown: 7 Apr. Weedkillers applied: 5 May. Combine harvested: 22 Aug.

GRAIN TONNES/HECTARE

***** Tables of means *****

FUNGCIDE [1]	NONE	TRIADIM	Mean
WEEDKLLR			
NONE	4.31	4.17	4.24
GLYPHOS	4.29	4.38	4.34
Mean	4.30	4.28	4.29
FUNGCIDE [2]	NONE	BENOMYL	Mean
WEEDKLLR			
NONE	4.25	4.24	4.24
GLYPHOS	4.31	4.36	4.34
Mean	4.28	4.30	4.29
FUNGCIDE [2]	NONE	BENOMYL	Mean
FUNGCIDE [1]			
NONE	4.24	4.37	4.30
TRIADIM	4.32	4.24	4.28
Mean	4.28	4.30	4.29
INSCTCDE	NONE	CHLORFEN	Mean
WEEDKLLR			
NONE	4.25	4.24	4.24
GLYPHOS	4.29	4.38	4.34
Mean	4.27	4.31	4.29
INSCTCDE	NONE	CHLORFEN	Mean
FUNGCIDE [1]			
NONE	4.24	4.37	4.30
TRIADIM	4.30	4.25	4.28
Mean	4.27	4.31	4.29
INSCTCDE	NONE	CHLORFEN	Mean
FUNGCIDE [2]			
NONE	4.24	4.32	4.28
BENOMYL	4.30	4.30	4.30
Mean	4.27	4.31	4.29
NEMACIDE	NONE	ALDICARB	Mean
WEEDKLLR			
NONE	4.05	4.44	4.24
GLYPHOS	4.21	4.47	4.34
Mean	4.13	4.45	4.29

88/R/CS/140

GRAIN TONNES/HECTARE

***** Tables of means *****

NEMACIDE FUNGicide [1]	NONE	ALDICARB	Mean
NONE	4.08	4.53	4.30
TRIADIM	4.18	4.38	4.28
Mean	4.13	4.45	4.29
NEMACIDE FUNGicide [2]	NONE	ALDICARB	Mean
NONE	4.16	4.40	4.28
BENOMYL	4.10	4.50	4.30
Mean	4.13	4.45	4.29
NEMACIDE INSCTCDE	NONE	ALDICARB	Mean
NONE	4.05	4.50	4.27
CHLORFEN	4.21	4.41	4.31
Mean	4.13	4.45	4.29
WEEDKLLR	FUNGicide [2] FUNGicide [1]	NONE	BENOMYL
NONE	NONE	4.30	4.33
	TRIADIM	4.20	4.15
GLYPHOS	NONE	4.19	4.40
	TRIADIM	4.43	4.33
WEEDKLLR	INSCTCDE FUNGicide [1]	NONE	CHLORFEN
NONE	NONE	4.24	4.38
	TRIADIM	4.26	4.09
GLYPHOS	NONE	4.24	4.35
	TRIADIM	4.34	4.41
WEEDKLLR	INSCTCDE FUNGicide [2]	NONE	CHLORFEN
NONE	NONE	4.26	4.24
	BENOMYL	4.24	4.23
GLYPHOS	NONE	4.22	4.39
	BENOMYL	4.36	4.37
FUNGicide [1]	INSCTCDE FUNGicide [2]	NONE	CHLORFEN
NONE	NONE	4.13	4.36
	BENOMYL	4.36	4.37
TRIADIM	NONE	4.36	4.27
	BENOMYL	4.24	4.23

88/R/CS/140

GRAIN TONNES/HECTARE

***** Tables of means *****

WEEDKLLR	FUNGCID[1]	NEMACIDE	NONE	ALDICARB
		NONE		
GLYPHOS	GLYPHOS	TRIADIM	4.06	4.29
		NONE	4.12	4.47
WEEDKLLR	FUNGCID[2]	TRIADIM	4.30	4.46
		NEMACIDE	NONE	ALDICARB
GLYPHOS	GLYPHOS	NONE	4.05	4.46
		BENOMYL	4.06	4.41
FUNGCID[1]	FUNGCID[2]	NONE	4.27	4.35
		BENOMYL	4.15	4.58
FUNGCID[1]	FUNGCID[2]	NEMACIDE	NONE	ALDICARB
		NONE	4.03	4.46
TRIADIM	TRIADIM	BENOMYL	4.14	4.59
		NONE	4.28	4.35
WEEDKLLR	INSCTCDE	BENOMYL	4.07	4.40
		NEMACIDE	NONE	ALDICARB
GLYPHOS	GLYPHOS	NONE	3.99	4.52
		CHLORFEN	4.12	4.36
FUNGCID[1]	INSCTCDE	NONE	4.11	4.47
		CHLORFEN	4.30	4.46
FUNGCID[1]	INSCTCDE	NEMACIDE	NONE	ALDICARB
		NONE	3.94	4.54
TRIADIM	TRIADIM	CHLORFEN	4.22	4.51
		NONE	4.16	4.45
FUNGCID[2]	INSCTCDE	CHLORFEN	4.20	4.31
		NEMACIDE	NONE	ALDICARB
BENOMYL	BENOMYL	NONE	4.02	4.47
		CHLORFEN	4.30	4.34
FUNGCID[2]	INSCTCDE	NONE	4.08	4.52
		CHLORFEN	4.12	4.48

88/R/CS/140

GRAIN TONNES/HECTARE

*** Standard errors of differences of means ***

Margins of two factor tables	0.063
Two factor tables	0.089
Three factor tables	0.126

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
WP	6	0.178	4.1

GRAIN MEAN DM% 85.2

PLOT AREA HARVESTED 0.00075