

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1989

[Full Table of Content](#)



### 89/R/CS/331 Take-all Inoculation - W. Wheat, W. Oats

#### Rothamsted Research

Rothamsted Research (1990) *89/R/CS/331 Take-all Inoculation - W. Wheat, W. Oats* ; Yields Of The Field Experiments 1989, pp 99 - 100 - DOI: <https://doi.org/10.23637/ERADOC-1-40>

89/R/CS/331

### TAKE-ALL INOCULATION

**Object:** To compare a range of methods of artificially inoculating take-all (*Gaeumannomyces graminis*) and to relate amounts of disease established to the yield and grain quality of w. wheat - Great Harpenden I.

**Sponsors:** D. Hornby, G.L. Bateman, R.J. Gutteridge.

The first year, w. wheat.

**Design:** 4 randomised blocks of 9 plots.

**Whole plot dimensions:** 3.0 x 22.0.

**Treatments:**

<b>INOCMETH</b>	Methods of inoculating take-all to w. wheat:
<b>NONE</b>	None (duplicated)
<b>I PRE PL</b>	Infective inoculum applied to soil surface pre-ploughing
<b>N PRE PL</b>	Non-infective inoculum applied to soil surface pre-ploughing
<b>I PRE SO</b>	Infective inoculum applied by fertilizer drill to 10 cm depth before rotary harrowing and sowing wheat
<b>N PRE SO</b>	Non-infective inoculum applied as above
<b>I CD</b>	Infective inoculum combine drilled with the seed
<b>N CD</b>	Non-infective inoculum combine drilled with the seed

**NOTES:** (1) Inoculum was prepared on autoclaved oat seed sown at 212 kg.  
(2) The sequence of cultivations was identical for all treatments: Plough to 23 cm on 5 Oct, 1988, cultivate to level on 22 Oct, traverse with fertilizer drill to 10 cm on 26 Oct, rotary harrow to 10 cm and sow wheat with combine drill on 27 Oct.  
(3) An additional treatment, required for comparisons in future years, was sown with w. oats.

**Basal applications:** Manures: (0:18:36) at 920 kg. 'Nitram' at 580 kg. Weedkillers: Paraquat at 0.60 kg ion in 200 l. Methabenzthiazuron at 1.6 kg in 200 l. Mecoprop at 2.2 kg, bromoxynil at 0.28 kg and ioxynil at 0.28 kg in 200 l. Fungicides: Prochloraz at 0.40 kg and carbendazim at 0.15 kg in 200 l.

**Seed:** W. wheat: Mercia, sown at 190 kg.  
W. oats: Image, sown at 190 kg.

**Cultivations, etc.:** - Rotary cultivated: 19 Aug, 1988. PK applied: 16 Sept. Paraquat applied: 19 Sept. Methabenzthiazuron applied: 3 Nov. N applied: 18 Apr, 1989. Mecoprop, bromoxynil and ioxynil applied: 4 May. Fungicides applied: 10 May. Combine harvested: 4 Aug (w. wheat), 17 Aug (w. oats). Previous crops: W. beans 1987, w. oats 1988.

**NOTE:** Plant samples were taken on six occasions between mid-March and mid-July to assess take-all. Quality assessments were made on the grain. Soil cores were taken after harvest to assess take-all infectivity, before and after cultivations.

89/R/CS/331

GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

INOCMETH	
NONE	7.11
I PRE PL	6.96
N PRE PL	7.11
I PRE SO	6.88
N PRE SO	7.35
I CD	6.58
N CD	7.12
Mean	7.03

\*\*\* Standard errors of differences of means \*\*\*

INOCMETH	
0.168	min.rep
0.145	max-min

INOCMETH	
max-min	NONE v any of the remainder
min.rep	any of the remainder

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	22	0.237	3.4
GRAIN MEAN DM%	87.1		
PLOT AREA HARVESTED	0.00506		