

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1989

[Full Table of Content](#)



89/R/CS/333 Comparison of Combinable Crops - W. Wheat

Rothamsted Research

Rothamsted Research (1990) *89/R/CS/333 Comparison of Combinable Crops - W. Wheat* ; Yields Of The Field Experiments 1989, pp 101 - 103 - DOI: <https://doi.org/10.23637/ERADOC-1-40>

89/R/CS/333

COMPARISON OF COMBINABLE CROPS

Object: To compare yields and other attributes of a range of combinable crops and to study their effects on a following crop of w. wheat - Long Hoos VI/VII 1.

Sponsors: J. McEwen, D.P. Yeoman, R.J. Darby, M.V. Hewitt.

The second year, w. wheat.

For previous year see 88/R/CS/333.

Design: 3 randomised blocks of 12 plots split into 2.

Whole plot dimensions: 2.5 x 8.0.

Treatments: All combinations of:-

Whole plots

1. PREVCROP	Crops in 1988:
W BEANS	W. beans, <i>Vicia faba</i>
W OATS	W. oats
W PEAS	W. peas, <i>Pisum sativum</i>
W RAPE	W. oilseed rape
W WHEAT	W. wheat
S BEANS	S. beans, <i>Vicia faba</i>
S LUPINS	S. lupins, <i>Lupinus albus</i>
S PEAS	S. peas, <i>Pisum sativum</i>
SNFLOWER	Sunflowers
FALLOW/C	Fallow, cultivated
FALLOW/P	Fallow, paraquat applied on four occasions
RYEGRASS	One-year ryegrass, cut and produce returned

Sub plots

2. SPRING N	Nitrogen fertilizer applied on 18 Apr, 1989:
O	None
N	N applied, amount depending on quantity in crop and soil in spring.

NOTE: Amounts of N applied (kg N) as 'Nitro-Chalk' were:

After PREVCROP	FALLOW/C	170
	W PEAS, S BEANS, S PEAS	180
	FALLOW/P	190
	W BEANS, W RAPE	200
	S LUPINS	210
	SUNFLOWERS	220
	W WHEAT, RYEGRASS	230
	W OATS	240

89/R/CS/333

Standard applications: Weedkillers: Paraquat at 0.60 kg ion in 220 l except after lupins. Isoproturon at 2.5 kg with mecoprop at 1.5 kg in 220 l except after lupins. Isoproturon at 2.5 kg with mecoprop at 0.42 kg in 220 l after lupins only. Isoproturon at 2.5 kg with mecoprop at 1.7 kg with cyanazine at 0.46 kg and clopyralid at 0.08 kg in 220 l to all plots. Difenzoquat at 0.99 kg applied with a wetting agent ('Agral' at 1.0 l) in 220 l. Fungicides: Propiconazole at 0.12 kg in 220 l. Carbendazim at 0.15 kg, maneb at 1.6 kg and tridemorph at 0.38 kg in 220 l. Molluscicide: Metaldehyde at 0.60 kg.

Seed: W. wheat: Mercia, sown at 200 kg.

Cultivations, etc.: - Paraquat applied except after lupins: 9 Sept, 1988. Deep-tine cultivated, except after lupins: 12 Sept. Rotary cultivated and seed sown except after lupins: 19 Sept. After lupin plots spring-tine cultivated, seed sown: 4 Oct. Isoproturon and mecoprop applied except after lupins: 19 Oct, and after lupins: 31 Oct and again to all plots with cyanazine and clopyralid: 6 Jan, 1989. Difenzoquat applied: 8 Feb. Propiconazole applied: 8 Mar. Metaldehyde applied: 15 Mar. Carbendazim, maneb and tridemorph applied: 26 Apr. Combine harvested: 28 July.

NOTE: Because the grain was spilt before weighing, the yield of one plot was lost, with treatment S BEANS N. An estimated value was used in the analysis.

89/R/CS/333

GRAIN TONNES/HECTARE

***** Tables of means *****

SPRING N PREVCROP	O	N	Mean
W BEANS	6.75	9.63	8.19
W OATS	3.82	8.81	6.31
W PEAS	6.82	9.17	7.99
W RAPE	6.61	9.24	7.92
W WHEAT	2.51	7.20	4.85
S BEANS	7.47	9.80	8.64
S LUPINS	6.66	9.42	8.04
S PEAS	6.96	9.51	8.24
SNFLOWER	4.86	9.14	7.00
FALLOW/C	8.10	9.70	8.90
FALLOW/P	7.43	9.75	8.59
RYEGRASS	3.68	7.55	5.62
Mean	5.97	9.08	7.52

*** Standard errors of differences of means ***

	PREVCROP	SPRING N	PREVCROP SPRING N
	0.380	0.157	0.541
Except when comparing means with the same level(s) of PREVCROP			0.545

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	22	0.465	6.2
BLOCK.WP.SP	23	0.667	8.9

GRAIN MEAN DM% 88.0

SUB PLOT AREA HARVESTED 0.00055