


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1989

[Full Table of Content](#)



### 89/R/WW/4 Factors Affecting Take-all - W. Wheat

#### Rothamsted Research

Rothamsted Research (1990) *89/R/WW/4 Factors Affecting Take-all - W. Wheat* ; Yields Of The Field Experiments 1989, pp 122 - 126 - DOI: <https://doi.org/10.23637/ERADOC-1-40>

89/R/WW/4

WINTER WHEAT

FACTORS AFFECTING TAKE-ALL

**Object:** To study the effects of a range of factors on the incidence of take-all and on the yield of w. wheat - White Horse II.

**Sponsors:** D. Hornby, G.L. Bateman, R.J. Gutteridge.

**Design:** A single replicate of 2 x 2 x 2 x 2 x 4.

**Whole plot dimensions:** 3.0 x 10.0.

**Treatments:** All combinations of:-

1. **SOWDATE**                      Dates of sowing:  
    22 SEPT                      22 September, 1988  
    28 OCT                        28 October
2. **SOILFUNG**                    Application of fungicide to the seedbed:  
    NONE                         None  
    NUARIMOL                    Nuarimol at 1.1 kg in 375 l
3. **SEEDRATE**                    Seed rates:  
    100 KG  
    200 KG
4. **AUTUMN N**                    N application to the seedbed:  
    0                              None  
    60                             60 kg N as 'Nitro-Chalk' on 21 Sept, 1988 or 27 Oct  
                                      for successive SOWDATES
5. **SPRING N**                    Nitrogen fertilizer (kg N) in spring, as 'Nitro-Chalk', applied 20 Apr, 1989:  
    100  
    150  
    200  
    250

**Basal applications:** Manures: (0:18:36) at 920 kg. Weedkillers: Chlortoluron at 3.5 kg in 200 l. Fluroxypyr at 0.20 kg in 200 l. Fungicide: Fenpropimorph at 0.75 kg in 200 l.

**Cultivations, etc.:-** Heavy spring-tine cultivated: 17 Sept, 1988. PK applied: 19 Sept. Rotary harrowed: 20 Sept. Chlortoluron applied: 16 Nov. Fluroxypyr applied: 8 May 1989. Fungicide applied: 23 May. Combine harvested: 8 Aug. Previous crops: W. wheat 1987 and 1988.

**NOTE:** Plant samples were taken in mid-April and June to assess take-all, eyespot, sharp eyespot, damage by stem-boring larvae (April only) and brown foot rot (June only). Grain quality was measured.

89/R/WW/4

GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

<b>SOILFUNG</b>	NONE	NUARIMOL	Mean		
<b>SOWDATE</b>					
22 SEPT	6.58	6.81	6.69		
28 OCT	6.91	6.99	6.95		
Mean	6.74	6.90	6.82		
<b>SEEDRATE</b>	100 KG	200 KG	Mean		
<b>SOWDATE</b>					
22 SEPT	6.68	6.70	6.69		
28 OCT	6.86	7.03	6.95		
Mean	6.77	6.87	6.82		
<b>SEEDRATE</b>	100 KG	200 KG	Mean		
<b>SOILFUNG</b>					
NONE	6.68	6.81	6.74		
NUARIMOL	6.86	6.93	6.90		
Mean	6.77	6.87	6.82		
<b>AUTUMN N</b>	0	60	Mean		
<b>SOWDATE</b>					
22 SEPT	6.55	6.84	6.69		
28 OCT	6.86	7.04	6.95		
Mean	6.70	6.94	6.82		
<b>AUTUMN N</b>	0	60	Mean		
<b>SOILFUNG</b>					
NONE	6.58	6.91	6.74		
NUARIMOL	6.82	6.97	6.90		
Mean	6.70	6.94	6.82		
<b>AUTUMN N</b>	0	60	Mean		
<b>SEEDRATE</b>					
100 KG	6.68	6.86	6.77		
200 KG	6.72	7.01	6.87		
Mean	6.70	6.94	6.82		
<b>SPRING N</b>	100	150	200	250	Mean
<b>SOWDATE</b>					
22 SEPT	6.52	6.79	6.76	6.71	6.69
28 OCT	6.88	7.01	6.99	6.92	6.95
Mean	6.70	6.90	6.87	6.81	6.82

89/R/WW/4

GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

<b>SPRING N</b>	100	150	200	250	Mean
<b>SOILFUNG</b>					
NONE	6.51	6.88	6.93	6.66	6.74
NUARIMOL	6.89	6.92	6.82	6.97	6.90
Mean	6.70	6.90	6.87	6.81	6.82
<b>SPRING N</b>	100	150	200	250	Mean
<b>SEEDRATE</b>					
100 KG	6.70	6.83	6.85	6.70	6.77
200 KG	6.70	6.96	6.90	6.93	6.87
Mean	6.70	6.90	6.87	6.81	6.82
<b>SPRING N</b>	100	150	200	250	Mean
<b>AUTUMN N</b>					
0	6.49	6.74	6.84	6.74	6.70
60	6.91	7.06	6.90	6.89	6.94
Mean	6.70	6.90	6.87	6.81	6.82
<b>SOWDATE</b>	<b>SEEDRATE</b>	100 KG	200 KG		
	<b>SOILFUNG</b>				
22 SEPT	NONE	6.53	6.63		
	NUARIMOL	6.83	6.78		
28 OCT	NONE	6.83	6.98		
	NUARIMOL	6.89	7.09		
<b>SOWDATE</b>	<b>AUTUMN N</b>	0	60		
	<b>SOILFUNG</b>				
22 SEPT	NONE	6.41	6.75		
	NUARIMOL	6.68	6.93		
28 OCT	NONE	6.76	7.06		
	NUARIMOL	6.96	7.02		
<b>SOWDATE</b>	<b>AUTUMN N</b>	0	60		
	<b>SEEDRATE</b>				
22 SEPT	100 KG	6.49	6.87		
	200 KG	6.60	6.81		
28 OCT	100 KG	6.87	6.85		
	200 KG	6.85	7.22		
<b>SOILFUNG</b>	<b>AUTUMN N</b>	0	60		
	<b>SEEDRATE</b>				
NONE	100 KG	6.50	6.87		
	200 KG	6.67	6.95		
NUARIMOL	100 KG	6.86	6.86		
	200 KG	6.78	7.08		

89/R/WW/4

GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

	<b>SPRING N</b>	100	150	200	250
<b>SOWDATE</b>	<b>SOILFUNG</b>				
22 SEPT	NONE	6.18	6.76	6.94	6.44
	NUARIMOL	6.87	6.81	6.57	6.98
28 OCT	NONE	6.85	6.99	6.91	6.88
	NUARIMOL	6.90	7.03	7.06	6.96
	<b>SPRING N</b>	100	150	200	250
<b>SOWDATE</b>	<b>SEEDRATE</b>				
22 SEPT	100 KG	6.56	6.71	6.87	6.58
	200 KG	6.48	6.86	6.64	6.83
28 OCT	100 KG	6.84	6.95	6.83	6.81
	200 KG	6.91	7.06	7.15	7.02
	<b>SPRING N</b>	100	150	200	250
<b>SOILFUNG</b>	<b>SEEDRATE</b>				
NONE	100 KG	6.48	6.92	6.81	6.52
	200 KG	6.55	6.83	7.05	6.79
NUARIMOL	100 KG	6.93	6.74	6.89	6.88
	200 KG	6.84	7.09	6.74	7.06
	<b>SPRING N</b>	100	150	200	250
<b>SOWDATE</b>	<b>AUTUMN N</b>				
22 SEPT	0	6.39	6.66	6.60	6.54
	60	6.66	6.91	6.91	6.88
28 OCT	0	6.58	6.82	7.08	6.94
	60	7.17	7.20	6.89	6.89
	<b>SPRING N</b>	100	150	200	250
<b>SOILFUNG</b>	<b>AUTUMN N</b>				
NONE	0	6.36	6.66	6.86	6.44
	60	6.67	7.09	6.99	6.88
NUARIMOL	0	6.61	6.81	6.82	7.04
	60	7.16	7.02	6.81	6.90
	<b>SPRING N</b>	100	150	200	250
<b>SEEDRATE</b>	<b>AUTUMN N</b>				
100 KG	0	6.44	6.72	6.88	6.67
	60	6.97	6.95	6.81	6.72
200 KG	0	6.53	6.76	6.80	6.81
	60	6.86	7.17	6.99	7.05

89/R/WW/4

GRAIN TONNES/HECTARE

\*\*\* Standard errors of differences of means \*\*\*

SOWDATE	SOILFUNG	SEEDRATE	AUTUMN N
0.064	0.064	0.064	0.064
SPRING N	SOWDATE	SOWDATE	SOILFUNG
	SOILFUNG	SEEDRATE	SEEDRATE
0.090	0.090	0.090	0.090
SOWDATE	SOILFUNG	SEEDRATE	SOWDATE
AUTUMN N	AUTUMN N	AUTUMN N	SPRING N
0.090	0.090	0.090	0.128
SOILFUNG	SEEDRATE	AUTUMN N	SOWDATE
SPRING N	SPRING N	SPRING N	SOILFUNG
			SEEDRATE
0.128	0.128	0.128	0.128
SOWDATE	SOWDATE	SOILFUNG	SOWDATE*
SOILFUNG	SEEDRATE	SEEDRATE	SOILFUNG
AUTUMN N	AUTUMN N	AUTUMN N	SPRING N
0.128	0.128	0.128	0.180
SOWDATE	SOILFUNG	SOWDATE	SOILFUNG
SEEDRATE	SEEDRATE	AUTUMN N	AUTUMN N
SPRING N	SPRING N	SPRING N	SPRING N
0.180	0.180	0.180	0.180
SEEDRATE			
AUTUMN N			
SPRING N			
0.180			

\* Only when comparing means with the same level of SOWDATE.SOILFUNG.NIT, where NIT has 2 levels. The first corresponding to levels 100, 250 of SPRING N and the second to the other 2 levels.

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	14	0.255	3.7
GRAIN MEAN DM%	89.1		
PLOT AREA HARVESTED	0.00226		