

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1989

[Full Table of Content](#)



---

### 89/R/B/2 Sowing Dates, Aphids and Bydv - W. Barley

#### Rothamsted Research

Rothamsted Research (1990) *89/R/B/2 Sowing Dates, Aphids and Bydv - W. Barley* ; Yields Of The Field Experiments 1989, pp 151 - 152 - DOI: <https://doi.org/10.23637/ERADOC-1-40>

89/R/B/2

WINTER BARLEY

SOWING DATES, APHIDS AND BYDV

**Object:** To study the relationship of aphid numbers in suction trap samples to crop populations and the incidence of BYDV on winter barley sown on a range of dates - Great Field II.

**Sponsors:** N. Carter, R.J. Gutteridge, J.F. Jenkyn, R.T. Plumb.

**Design:** 4 randomised blocks of 10 plots.

**Whole plot dimensions:** 3.0 x 23.0 N.BLOCKS.  
3.0 x 18.0 S.BLOCKS.

**Treatments:** All combinations of:-

1. **SOWDATE**                      Dates of sowing:  
  
    12 SEPT                      12 September, 1988  
    22 SEPT                      22 September  
    3 OCT                        3 October  
    17 OCT                      17 October  
    27 OCT                      27 October
  
2. **APHICIDE**                    Aphicide:  
  
    NONE                        None  
    CYPERMET                  Cypermethrin at 0.025 kg in 380 l on 7 Nov, 1988  
                                      except on SOWDATE 27 OCT, applied on 11 Jan, 1989

**NOTE:** All SOWDATE treatments were cultivated by rotary grubber on 12 Sept, 1988 and rotary harrowed on the day of sowing.

**Basal applications:** Manures: 'Nitram' at 480 kg. Weedkillers: Glyphosate at 0.27 kg in 200 l. Chlortoluron at 3.5 kg in 200 l. Isoproturon at 1.5 kg with mecoprop at 2.2 kg, ioxynil at 0.28 kg and bromoxynil at 0.28 kg in 200 l. Fungicides: Propiconazole at 0.12 kg with tridemorph at 0.38 kg in 200 l.

**Seed:** Igri, sown at 150 kg.

**Cultivations, etc.:-** Rotary cultivated: 11 Aug, 1988, and 15 Aug. Glyphosate applied: 6 Sept. Chlortoluron applied: 16 Nov. N applied: 19 Apr, 1989. Remaining weedkillers applied: 2 May. Fungicides applied: 10 May. Combine harvested: 14 July. Previous crops: W. barley 1987 and 1988.

**NOTE:** Aphids were sampled from late September to June. Visual estimates of BYDV were made throughout the season, a more detailed estimate was made at the end of April. Components of yield were measured. Take-all was assessed in summer.

89/R/B/2

GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

APHICIDE SOWDATE	NONE	CYPERMET	Mean
12 SEPT	3.22	3.79	3.50
22 SEPT	3.54	4.26	3.90
3 OCT	3.49	3.75	3.62
17 OCT	2.92	3.37	3.14
27 OCT	3.42	3.68	3.55
Mean	3.32	3.77	3.54

\*\*\* Standard errors of differences of means \*\*\*

SOWDATE	APHICIDE	SOWDATE APHICIDE
0.166	0.105	0.235

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
BLOCK.WP	27	0.332	9.4
GRAIN MEAN DM%	85.0		
PLOT AREA HARVESTED	0.00230		