

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1989

[Full Table of Content](#)



89/R/LP/1 Varieties, Sowing Dates and Seed Dressings - W. Lupins

Rothamsted Research

Rothamsted Research (1990) *89/R/LP/1 Varieties, Sowing Dates and Seed Dressings - W. Lupins* ; Yields Of The Field Experiments 1989, pp 162 - 164 - DOI:

<https://doi.org/10.23637/ERADOC-1-40>

89/R/LP/1

WINTER LUPINS

VARIETIES, SOWING DATES AND SEED DRESSINGS

Object: To compare three varieties of lupin (*Lupinus albus*) sown at two sowing dates with and without a seed dressing - Long Hoos III 6.

Sponsors: J. McEwen, H.L. Jones, D.P. Yeoman, A.W. Ferguson, J.F. Jenkyn.

Design: 2 randomised blocks of 12 plots.

Whole plot dimensions: 1.8 x 9.0.

Treatments: All combinations of:-

Whole plots

- | | |
|---------------------|--|
| 1. VARIETY | Varieties: |
| C 8 | C 8 |
| LUCKY | Lucky |
| VLADIMIR | Vladimir |
| 2. SOW DATE | Dates of sowing: |
| 21 SEPT | 21 September, 1988 |
| 2 NOV | 2 November |
| 3. FUNGICIDE | Fungicides: |
| NONE | None |
| F+C+T+B | Fosetyl-aluminium, captan and thiabendazole, to seed plus benomyl at 0.50 kg and chlorothalonil at 1.0 kg applied in 220 l on 15 Mar, 1989 |

NOTE: Plots were netted from October to May.

Basal applications: Manure: Muriate of potash at 520 kg. Weedkillers: Terbutryne at 0.98 kg and terbuthylazine at 0.42 kg in 220 l repeated at 0.49 kg and 0.21 kg respectively in 220 l to first sowing only. Metamitron at 2.8 kg applied with the benomyl, chlorothalonil and pirimicarb in 220 l. Fungicides: Benomyl at 0.50 kg with chlorothalonil at 1.0 kg. Benomyl at 0.56 kg applied with pirimicarb in 220 l. Insecticides: Pirimicarb at 0.14 kg on two occasions. Deltamethrin at 7.5 g in 220 l. Molluscicides: Methiocarb at 0.60 kg to first sowing and at 0.27 kg to second sowing. Metaldehyde at 0.94 kg.

Seed: Sown at 60 seeds per square metre.

89/R/LP/1

Cultivations, etc.:- Ploughed: 7 June, 1988. Rotary cultivated: 5 Aug. K applied: 15 Sept. Rotary harrowed, first sowing date seed sown: 21 Sept. Terbutryne and terbuthylazine applied: 29 and 30 Sept to first sowing only. Methiocarb applied to first sowing: 26 Oct. Second sowing date seed sown, terbutryne and terbuthylazine applied and methiocarb applied: 2 Nov. Metaldehyde applied: 15 Mar, 1989. Metamitron, benomyl, chlorothalonil and pirimicarb applied: 28 Apr. Deltamethrin applied: 9 May. Pirimicarb and benomyl applied: 9 June. Combine harvested: 23 Aug. Previous crops : Potatoes 1987, fallow 1988.

NOTE: All plots sown with **VARIETY VLADIMIR** failed and have been omitted from the analysis.

89/R/LP/1

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

FUNGCIDE	NONE	F+C+T+B	Mean
VARIETY			
C 8	2.10	1.31	1.71
LUCKY	2.08	1.75	1.91
Mean	2.09	1.53	1.81
SOW DATE 21 SEPT 2 NOV Mean			
VARIETY			
C 8	1.82	1.59	1.71
LUCKY	1.76	2.07	1.91
Mean	1.79	1.83	1.81
SOW DATE 21 SEPT 2 NOV Mean			
FUNGCIDE			
NONE	2.21	1.98	2.09
F+C+T+B	1.37	1.68	1.53
Mean	1.79	1.83	1.81
VARIETY SOW DATE 21 SEPT 2 NOV			
FUNGCIDE			
C 8	NONE	2.29	1.92
	F+C+T+B	1.36	1.27
LUCKY	NONE	2.13	2.04
	F+C+T+B	1.39	2.10

*** Standard errors of differences of means ***

VARIETY	FUNGCIDE	SOW DATE	VARIETY FUNGCIDE
0.202	0.202	0.202	0.286
VARIETY	FUNGCIDE	VARIETY	
SOW DATE	SOW DATE	FUNGCIDE	
		SOW DATE	
0.286	0.286	0.404	

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	7	0.404	22.3
GRAIN MEAN DM%	87.7		
PLOT AREA HARVESTED	0.00162		