


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1989

[Full Table of Content](#)



Lupins

Rothamsted Research

Rothamsted Research (1990) *Lupins* ; Yields Of The Field Experiments 1989, pp 162 - 166 - DOI: <https://doi.org/10.23637/ERADOC-1-40>

89/R/LP/1

WINTER LUPINS

VARIETIES, SOWING DATES AND SEED DRESSINGS

Object: To compare three varieties of lupin (*Lupinus albus*) sown at two sowing dates with and without a seed dressing - Long Hoos III 6.

Sponsors: J. McEwen, H.L. Jones, D.P. Yeoman, A.W. Ferguson, J.F. Jenkyn.

Design: 2 randomised blocks of 12 plots.

Whole plot dimensions: 1.8 x 9.0.

Treatments: All combinations of:-

Whole plots

- | | |
|---------------------|--------------------------------------------------------------------------------------------------------------------------------------------|
| 1. VARIETY | Varieties: |
| C 8 | C 8 |
| LUCKY | Lucky |
| VLADIMIR | Vladimir |
| 2. SOW DATE | Dates of sowing: |
| 21 SEPT | 21 September, 1988 |
| 2 NOV | 2 November |
| 3. FUNGICIDE | Fungicides: |
| NONE | None |
| F+C+T+B | Fosetyl-aluminium, captan and thiabendazole, to seed plus benomyl at 0.50 kg and chlorothalonil at 1.0 kg applied in 220 l on 15 Mar, 1989 |

NOTE: Plots were netted from October to May.

Basal applications: Manure: Muriate of potash at 520 kg. Weedkillers: Terbutryne at 0.98 kg and terbuthylazine at 0.42 kg in 220 l repeated at 0.49 kg and 0.21 kg respectively in 220 l to first sowing only. Metamitron at 2.8 kg applied with the benomyl, chlorothalonil and pirimicarb in 220 l. Fungicides: Benomyl at 0.50 kg with chlorothalonil at 1.0 kg. Benomyl at 0.56 kg applied with pirimicarb in 220 l. Insecticides: Pirimicarb at 0.14 kg on two occasions. Deltamethrin at 7.5 g in 220 l. Molluscicides: Methiocarb at 0.60 kg to first sowing and at 0.27 kg to second sowing. Metaldehyde at 0.94 kg.

Seed: Sown at 60 seeds per square metre.

89/R/LP/1

Cultivations, etc.:- Ploughed: 7 June, 1988. Rotary cultivated: 5 Aug. K applied: 15 Sept. Rotary harrowed, first sowing date seed sown: 21 Sept. Terbutryne and terbutylazine applied: 29 and 30 Sept to first sowing only. Methiocarb applied to first sowing: 26 Oct. Second sowing date seed sown, terbutryne and terbutylazine applied and methiocarb applied: 2 Nov. Metaldehyde applied: 15 Mar, 1989. Metamitron, benomyl, chlorothalonil and pirimicarb applied: 28 Apr. Deltamethrin applied: 9 May. Pirimicarb and benomyl applied: 9 June. Combine harvested: 23 Aug. Previous crops : Potatoes 1987, fallow 1988.

NOTE: All plots sown with **VARIETY VLADIMIR** failed and have been omitted from the analysis.

89/R/LP/1

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

FUNGCIDE	NONE	F+C+T+B	Mean
VARIETY			
C 8	2.10	1.31	1.71
LUCKY	2.08	1.75	1.91
Mean	2.09	1.53	1.81
SOW DATE	21 SEPT	2 NOV	Mean
VARIETY			
C 8	1.82	1.59	1.71
LUCKY	1.76	2.07	1.91
Mean	1.79	1.83	1.81
SOW DATE	21 SEPT	2 NOV	Mean
FUNGCIDE			
NONE	2.21	1.98	2.09
F+C+T+B	1.37	1.68	1.53
Mean	1.79	1.83	1.81
VARIETY	SOW DATE	21 SEPT	2 NOV
	FUNGCIDE		
C 8	NONE	2.29	1.92
	F+C+T+B	1.36	1.27
LUCKY	NONE	2.13	2.04
	F+C+T+B	1.39	2.10

*** Standard errors of differences of means ***

VARIETY	FUNGCIDE	SOW DATE	VARIETY FUNGCIDE
0.202	0.202	0.202	0.286
VARIETY	FUNGCIDE	VARIETY	
SOW DATE	SOW DATE	FUNGCIDE	
SOW DATE		SOW DATE	
0.286	0.286	0.404	

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	7	0.404	22.3
GRAIN MEAN DM%	87.7		
PLOT AREA HARVESTED	0.00162		

89/R/LP/2

SPRING LUPINS

VERNALIZATION STUDY

Object: To study the effects of a range of periods of vernalization on the growth, date of maturity and yield of two varieties of lupins (*Lupinus albus*) - Long Hoos IV O & E S.W.

Sponsors: G.F.J. Milford, L.A. Mullen, D.P. Yeoman.

Design: A single replicate of 28 plots.

Whole plot dimensions: 1.5 x 2.0.

Treatments: All combinations of:-

1. **SOWDATE** Dates of sowing:

8 MARCH
29 MARCH

2. **VARIETY** Varieties:

C8
VLADIMIR

3. **VERNTIME** Period of vernalization at 1 degree C:

NONE
1 WEEK
2 WEEKS
3 WEEKS
4 WEEKS
5 WEEKS
6 WEEKS

Basal applications: Manures: (0:18:36) at 1080 kg. Magnesian limestone at 2.6 t. Weedkillers: Glyphosate at 0.72 kg in 220 l. Paraquat at 0.54 kg ion in 220 l. Terbutryne at 0.98 kg and terbuthylazine at 0.42 kg in 220 l. Insecticides: Deltamethrin at 7.5 g in 220 l. Pirimicarb at 0.14 kg in 220 l. Molluscicides: Methiocarb at 0.40 kg applied twice. Metaldehyde at 0.94 kg.

Cultivations, etc.:- Glyphosate applied: 13 Oct, 1988. PK applied: 26 Oct. Paraquat applied: 24 Nov. Magnesian limestone applied: 30 Nov. Ploughed: 8 Dec. Rotary cultivated, first sowing date seed sown, methiocarb applied: 8 Mar, 1989. Terbutryne and terbuthylazine applied to first sowing, methiocarb applied: 15 Mar. Rotary cultivated, second sowing date seed sown, rolled: 29 Mar. Terbutryne and terbuthylazine applied to second sowing: 30 Mar. Metaldehyde applied: 31 Mar. Deltamethrin applied: 9 May. Pirimicarb applied: 5 July. Hand harvested: 20 Sept. Previous crops: S.barley 1987 and 1988.

89/R/LP/2

NOTES: (1) All plots sown with **VARIETY** VLADIMIR failed and only the results from **VARIETY** C8 are presented.
(2) Grain mean dry matter percentages are not available.

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

SOW DATE	8 MARCH	29 MARCH	Mean
VERTIME			
NONE	0.80	0.93	0.86
1 WEEK	0.42	0.59	0.51
2 WEEKS	0.09	0.92	0.51
3 WEEKS	0.81	1.14	0.97
4 WEEKS	0.53	0.85	0.69
5 WEEKS	0.88	1.29	1.08
6 WEEKS	0.21	1.14	0.67
Mean	0.53	0.98	0.76

PLOT AREA HARVESTED 0.00030