Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1989



Full Table of Content

89/R/RA/3 Varieties, Pests and Diseases - W. Oilseed Rape

Rothamsted Research

Rothamsted Research (1990) 89/R/RA/3 Varieties, Pests and Diseases - W. Oilseed Rape; Yields Of The Field Experiments 1989, pp 180 - 182 - DOI: https://doi.org/10.23637/ERADOC-1-40

89/R/RA/3

WINTER OILSEED RAPE

VARIETIES, PESTS AND DISEASES

Object: To investigate the effects of full pest and disease control on a range of low glucosinolate varieties compared with Bienvenu - Osier.

Sponsors: C.J. Rawlinson, I.H. Williams.

Design: 2 randomised blocks of 24 plots.

Whole plot dimensions: 3.0 x 21.0.

Treatments: All combinations of:-

VARIETY Varieties:

ARIANA Ariana
BIENVENU Bienvenu
CAPRCORN Capricorn
COBRA Cobra
LIBRAVO Libravo
TAPIDOR Tapidor

2. INSCTCDE Insecticides:

NONE None

FULL Deltamethrin at 6.2 g in 200 l on 1 Oct, 1988, and 7 Nov. Azinphos methyl at 0.40 kg and demeton-Smethyl sulphone at 0.12 kg in 300 l on 29 Mar, 1989. Triazophos at 0.42 kg in 260 l on 2 June.

3. FUNGCIDE Fungicides:

NONE None

FULL Prochloraz at 0.50 kg in 200 1 on 7 Nov, 1988 and 29 Mar, 1989. Iprodione at 0.50 kg in 260 1 on 2 June.

Basal applications: Manure: 'Nitram' at 140 kg on two occasions and on
 a third occasion at 290 kg. Weedkillers: TCA at 10 kg in 200 l.
 Metazachlor at 1.2 kg with fluazifop-p-butyl at 0.19 kg and a wetting
 agent ('Enhance' at 0.26 l) in 260 l. Desiccant: Diquat at 0.60 kg
 ion with a wetting agent ('Enhance' at 0.50 l) in 520 l.

Seed: Varieties, sown at 8.0 kg.

Cultivations, etc.:- Heavy spring-tine cultivated, rotary cultivated:
9 Aug, 1988. First N applied, TCA applied: 16 Aug. Seed sown:
22 Aug. Remaining weedkillers with wetting agent applied: 30 Sept.
Second N applied: 2 Mar, 1989. Third N applied: 28 Mar. Desiccant with wetting agent applied: 12 July. Combine harvested: 15 July.
Previous crops: W. wheat 1987, w. barley 1988.

89/R/RA/3

NOTE: Disease assessments were made on eight occasions throughout the season and pest numbers were recorded in autumn, spring and summer. Microflora of leaf and pods were assessed until harvest. Glucosinolate contents in plant tissues and grain were measured throughout the season.

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

**** Tables of means ****

INSCTCDE	NONE	FULL	Mean
VARIETY			
ARIANA	2.71	3.26	2.99
BIENVENU	3.39	3.41	3.40
CAPRCORN	2.87	4.02	3.45
COBRA	2.64	3.46	3.05
LIBRAVO	2.85	3.52	3.18
TAPIDOR	2.70	3.63	3.17
Mean	2.86	3.55	3.21
FUNGCIDE	NONE	FULL	Mean
VARIETY			
ARIANA	2.73	3.24	2.99
BIENVENU	3.24	3.56	3.40
CAPRCORN	3.29	3.61	3.45
COBRA	2.95	3.14	3.05
LIBRAVO	3.10	3.27	3.18
TAPIDOR	3.03	3.30	3.17
Mean	3.06	3.35	3.21
FUNGCIDE	NONE	FULL	Mean
INSCTCDE			
NONE	2.73	2.99	2.86
FULL	3.38	3.72	3.55
Mean	3.06	3.35	3.21
	FUNGCIDE	NONE	FULL
VARIETY	INSCTCDE		
ARIANA	NONE	2.58	2.84
	FULL	2.88	3.65
BIENVENU	NONE	3.36	3.42
	FULL	3.11	3.70
CAPRCORN	NONE	2.54	3.20
	FULL	4.03	4.01
COBRA	NONE	2.58	2.70
	FULL	3.33	3.59
LIBRAVO	NONE	2.78	2.93
	FULL	3.43	3.61
TAPIDOR	NONE	2.54	2.86
	FULL	3.53	3.74

89/R/RA/3

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

*** Standard errors of differences of means ***

VARIETY	INSCTCDE	FUNGCIDE	VARIETY INSCTCDE
0.163	0.094	0.094	0.230
VARIETY	INSCTCDE	VARIETY	
FUNGCIDE	FUNGCIDE	INSCTCDE	
		FUNGCIDE	
0.230	0.133	0.325	

***** Stratum standard errors and coefficients of variation ****

 Stratum
 d.f.
 s.e.
 cv%

 BLOCK.WP
 23
 0.325
 10.2

GRAIN MEAN DM% 78.0

PLOT AREA HARVESTED 0.00405