

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1989

[Full Table of Content](#)



89/R/CS/140 Chemical Reference Plots - S. Barley

Rothamsted Research

Rothamsted Research (1990) *89/R/CS/140 Chemical Reference Plots - S. Barley* ; Yields Of The Field Experiments 1989, pp 71 - 74 - DOI: <https://doi.org/10.23637/ERADOC-1-40>

89/R/CS/140

CHEMICAL REFERENCE PLOTS

Object: To study the persistence in soil of agricultural chemicals applied annually, singly and in combination and their effects on soil microflora and on yield of continuous s. barley - Long Hoos V 3.

Sponsors: R.H. Bromilow, A.A. Evans, P.H. Nicholls.

The 16th year, s. barley.

For previous years see 74-88/R/CS/140.

Design: Single replicate of 32 plots.

Whole plot dimensions: 4.06 x 4.57.

Treatments, applied cumulatively every year except as stated:

All combinations of:-

1. **WEEDKLLR** Weedkiller in autumn:
NONE None
GLYPHOS Glyphosate at 1.4 kg to barley stubble each autumn from 1979 to 1984, at 0.72 kg in 1985, at 0.54 kg in 1986, at 1.3 kg in 1987 and at 1.5 kg in 1988.
2. **FUNGICIDE[1]** Fungicide in autumn:
NONE None
TRIADIM Triadimefon at 0.25 kg in autumn 1981, 1982, 1984, 1985, 1986, 1987 and 1988, 0.28 kg in autumn 1983
3. **FUNGICIDE[2]** Fungicide in spring:
NONE None
BENOMYL Benomyl at 4 kg to seedbed
4. **INSCTCDE** Insecticide:
NONE None
CHLORFEN Chlorfenvinphos at 2 kg to the seedbed
5. **NEMACIDE** Nematicide:
NONE None
ALDICARB Aldicarb at 6 kg to the seedbed

NOTE: Glyphosate and triadimefon were applied in 220 1 on 10 Oct, 1988 and 24 Oct respectively. Other treatments were applied on 31 Mar, 1989.

Basal applications: Manure: 'Nitram' at 320 kg. Weedkillers: Bentazone at 0.80 kg, dichlorprop at 1.1 kg and MCPA at 0.64 kg in 220 1.

Seed: Klaxon, seed not dressed, sown at 160 kg.

89/R/CS/140

Cultivations, etc.:- Ploughed: 25 Nov, 1988. N applied: 7 Mar, 1989.
 Spring-tine cultivated, seedbed treatments applied, rotary harrowed,
 seed sown, rolled: 31 Mar. Weedkillers applied: 9 May. Combine
 harvested: 15 Aug.

GRAIN TONNES/HECTARE

***** Tables of means *****

FUNGCIDE [1]	NONE	TRIADIM	Mean
WEEDKLLR			
NONE	3.81	3.93	3.87
GLYPHOS	3.94	3.92	3.93
Mean	3.88	3.93	3.90

FUNGCIDE [2]	NONE	BENOMYL	Mean
WEEDKLLR			
NONE	3.85	3.89	3.87
GLYPHOS	3.90	3.97	3.93
Mean	3.87	3.93	3.90

FUNGCIDE [2]	NONE	BENOMYL	Mean
FUNGCIDE [1]			
NONE	3.83	3.93	3.88
TRIADIM	3.92	3.93	3.93
Mean	3.87	3.93	3.90

INSCTCDE	NONE	CHLORFEN	Mean
WEEDKLLR			
NONE	3.74	3.99	3.87
GLYPHOS	3.89	3.98	3.93
Mean	3.81	3.99	3.90

INSCTCDE	NONE	CHLORFEN	Mean
FUNGCIDE [1]			
NONE	3.72	4.03	3.88
TRIADIM	3.91	3.95	3.93
Mean	3.81	3.99	3.90

INSCTCDE	NONE	CHLORFEN	Mean
FUNGCIDE [2]			
NONE	3.80	3.94	3.87
BENOMYL	3.83	4.03	3.93
Mean	3.81	3.99	3.90

NEMACIDE	NONE	ALDICARB	Mean
WEEDKLLR			
NONE	3.63	4.10	3.87
GLYPHOS	3.73	4.14	3.93
Mean	3.68	4.12	3.90

89/R/CS/140

GRAIN TONNES/HECTARE

***** Tables of means *****

NEMACIDE	NONE	ALDICARB	Mean
FUNGCIDE [1]			
NONE	3.61	4.14	3.88
TRIADIM	3.75	4.10	3.93
Mean	3.68	4.12	3.90
NEMACIDE	NONE	ALDICARB	Mean
FUNGCIDE [2]			
NONE	3.67	4.08	3.87
BENOMYL	3.69	4.17	3.93
Mean	3.68	4.12	3.90
NEMACIDE	NONE	ALDICARB	Mean
INSCTCDE			
NONE	3.58	4.05	3.81
CHLORFEN	3.78	4.20	3.99
Mean	3.68	4.12	3.90
WEEDKLLR	FUNGCIDE [2]	NONE	BENOMYL
	FUNGCIDE [1]		
NONE	NONE	3.83	3.79
	TRIADIM	3.87	3.98
GLYPHOS	NONE	3.82	4.06
	TRIADIM	3.97	3.88
WEEDKLLR	INSCTCDE	NONE	CHLORFEN
	FUNGCIDE [1]		
NONE	NONE	3.63	3.99
	TRIADIM	3.85	4.00
GLYPHOS	NONE	3.81	4.07
	TRIADIM	3.96	3.89
WEEDKLLR	INSCTCDE	NONE	CHLORFEN
	FUNGCIDE [2]		
NONE	NONE	3.74	3.96
	BENOMYL	3.75	4.02
GLYPHOS	NONE	3.87	3.92
	BENOMYL	3.90	4.04
FUNGCIDE [1]	INSCTCDE	NONE	CHLORFEN
	FUNGCIDE [2]		
NONE	NONE	3.71	3.94
	BENOMYL	3.73	4.12
TRIADIM	NONE	3.89	3.95
	BENOMYL	3.92	3.95

89/R/CS/140

GRAIN TONNES/HECTARE

***** Tables of means *****

WEEDKLLR	FUNGCIDE [1]	NEMACIDE	NONE	ALDICARB
NONE	NONE		3.67	3.95
	TRIADIM		3.60	4.25
GLYPHOS	NONE		3.56	4.33
	TRIADIM		3.89	3.96

WEEDKLLR	FUNGCIDE [2]	NEMACIDE	NONE	ALDICARB
NONE	NONE		3.61	4.09
	BENOMYL		3.66	4.11
GLYPHOS	NONE		3.73	4.06
	BENOMYL		3.72	4.22

FUNGCIDE [1]	FUNGCIDE [2]	NEMACIDE	NONE	ALDICARB
NONE	NONE		3.61	4.04
	BENOMYL		3.61	4.24
TRIADIM	NONE		3.73	4.11
	BENOMYL		3.76	4.10

WEEDKLLR	INSCTCDE	NEMACIDE	NONE	ALDICARB
NONE	NONE		3.51	3.98
	CHLORFEN		3.76	4.22
GLYPHOS	NONE		3.66	4.11
	CHLORFEN		3.79	4.17

FUNGCIDE [1]	INSCTCDE	NEMACIDE	NONE	ALDICARB
NONE	NONE		3.55	3.89
	CHLORFEN		3.67	4.39
TRIADIM	NONE		3.61	4.20
	CHLORFEN		3.88	4.01

FUNGCIDE [2]	INSCTCDE	NEMACIDE	NONE	ALDICARB
NONE	NONE		3.47	4.14
	CHLORFEN		3.87	4.01
BENOMYL	NONE		3.69	3.96
	CHLORFEN		3.68	4.38

*** Standard errors of differences of means ***

Margins of two factor tables	0.144
Two factor tables	0.204
Three factor tables	0.289

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
WP	6	0.409	10.5

GRAIN MEAN DM% 84.6 PLOT AREA HARVESTED 0.00093