Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1990



Full Table of Content

90/R/WW/2 Control of Volunteers - W. Wheat

Rothamsted Research

Rothamsted Research (1991) 90/R/WW/2 Control of Volunteers - W. Wheat; Yields Of The Field Experiments 1990, pp 106 - 110 - DOI: https://doi.org/10.23637/ERADOC-1-42

90/R/WW/2

WINTER WHEAT

CONTROL OF VOLUNTEERS

Object: To compare methods of volunteer control in winter wheat after w. and s. barley - Delafield, Fosters Corner.

Sponsors: R. Moffitt, D.G. Christian.

Design: 3 replicates of 8 x 3 criss-cross.

Column plot dimensions: 6.0 x 20.0.

Treatments: All combinations of:-

Column plots

1. PRIMCULT Primary cultivations:

NONE None until just before sowing

DYNDRIVE 'Bomford Dynadrive'

PLOUGH Plough TINE Tine

2. CULTDATE Date of cultivations:

EARLY 25 July, 1989 (Delafield)

17 Aug (Fosters Corner)

LATER 15 Aug (Delafield)

7 Sept (Fosters Corner)

Row plots

3. PRSOWCON Pre-sowing volunteer control:

GLYPHOS Glyphosate at 0.27 kg in 200 l on 18 Oct, 1989

(Delafield), 1 Nov and 13 Nov (Fosters Corner)

PARAQUAT Paraquat at 0.60 kg ion in 200 l on 18 Oct

(Delafield), 1 Nov (Fosters Corner)

NONE None

- NOTES: (1) The 'Bomford Dynadrive' has a frame similar to a rotary cultivator but it has two rotating shafts containing flat, slightly twisted, spade-shaped tines. The front shaft drives the rear, it is fitted with twice the number of blades and rotates at about one third the speed of the rear shaft.
 - (2) A 1 m strip of Plaisant w. barley was broadcast on the surface, by drill at 120 kg on one end of each plot. It was broadcast before any treatments were applied on 25 July, 1989 (Delafield) and 17 Aug (Fosters Corner).
 - (3) All plots were heavy spring-tine cultivated on 1 Nov, 1989 (Delafield), 14 Nov (Fosters Corner) then rotary harrowed and seed sown by drill 7 Nov (Delafield), 15 Nov (Fosters Corner). The seed was spring-tine cultivated in, (Delafield), and harrowed in, (Fosters Corner), after sowing.
 - (4) PRIMCULT TINE was heavy spring-tine cultivated twice.

90/R/WW/2

Basal applications: Manures: 'Nitram' at 120 kg and later at 460 kg. Weedkillers: Chlorotoluron at 3.5 kg in 200 l. Bromoxynil at 0.25 kg, ioxynil at 0.25 kg and mecoprop at 2.0 kg with the prochloraz in 200 l (Delafield). Bromoxynil at 0.28 kg, ioxynil at 0.28 kg and mecoprop at 2.2 kg with the prochloraz in 200 l (Fosters Corner). Fungicides: Prochloraz at 0.40 kg. Chlorothalonil at 0.50 kg with propiconazole at 0.12 kg in 200 l (Delafield). Fenpropimorph at 0.75 kg in 200 l.

Seed: Mercia, sown at 180 kg.

Cultivations, etc.:- Chlorotoluron applied: 21 Nov, 1989 (Fosters Corner),
23 Nov (Delafield). N applied: 9 Mar, 1990 and 12 Apr. Bromoxynil,
ioxynil and mecoprop with prochloraz applied: 10 Apr (Delafield),
25 Apr (Fosters Corner). Chlorothalonil applied with the propiconazole
(Fosters Corner): 31 May. Fenpropimorph applied: 29 June. Combine
harvested: 10 Aug (Delafield), 13 Aug (Fosters Corner). Previous
crops: W. oilseed rape 1988, w. barley 1989 (Delafield); sunflowers
1988, s. barley 1989 (Fosters Corner).

NOTES: (1) Ears of volunteer plants were counted at anthesis of the sown crop.

(2) Percentage contamination of harvested grain by volunteer grain was measured.

90/R/WW/2 DELAFIELD W.WHEAT AFTER W.BARLEY

GRAIN TONNES/HECTARE

**** Tables of means ****

CULTDATE	EARLY	LATER	Mean	
PRIMCULT				
NONE	5.73	5.83	5.78	
DYNDRIVE	5.48	5.80	5.64	
PLOUGH	6.30	6.43	6.36	
TINE	5.84	5.89	5.86	
Mean	5.84	5.99	5.91	
PRSOWCON	GLYPHOS	PARAQUAT	NONE	Mean
PRIMCULT				
NONE	6.11	5.79	5.44	5.78
DYNDRIVE	5.85	5.94	5.14	5.64
PLOUGH	6.76	6.38	5.95	6.36
TINE	6.08	6.24	5.27	5.86
Mean	6.20	6.09	5.45	5.91
PRSOWCON	GLYPHOS	PARAQUAT	NONE	Mean
CULTDATE				
EARLY	6.10	5.92	5.50	5.84
LATER	6.30	6.25	5.40	5.99
Mean	6.20	6.09	5.45	5.91

90/R/WW/2 DELAFIELD W.WHEAT AFTER W.BARLEY

GRAIN TONNES/HECTARE

**** Tables of means ****

PRIMCULT	PRSOWCON	GLYPHOS	PARAQUAT	NONE
NONE	EARLY	6.18	5.57	5.44
	LATER	6.05	6.00	5.44
DYNDRIVE	EARLY	5.75	5.70	4.99
	LATER	5.94	6.17	5.29
PLOUGH	EARLY	6.61	6.22	6.08
	LATER	6.92	6.54	5.81
TINE	EARLY	5.85	6.18	5.48
	LATER	6.30	6.31	5.06

*** Standard errors of differences of means ***

	PRIMCULT	CULTDATE	PRSOWCON	PRIMCULT
	0.167	0.118	0.296	0.236
	PRIMCULT PRSOWCON	CULTDATE	PRIMCULT CULTDATE PRSOWCON	and species
	0.369	0.325	0.445	
Except when PRIMCULT	comparing means 0.354	with the same	level(s)	of
CULTDATE		0.317		
PRSOWCON	0.248	0.175	0.350	
PRIMCULT.CO	JLTDATE		0.419	

**** Stratum standard errors and coefficients of variation ****

Stratum	d.f.	s.e.	cv%
BLOCK.WP1	4	0.363	6.1
BLOCK.WP2	14	0.288	4.9
BLOCK.WP1.WP2	28	0.388	6.6

GRAIN MEAN DM% 89.5

PLOT AREA HARVESTED 0.00075

90/R/WW/2 FOSTERS CORNER W.WHEAT AFTER S.BARLEY

GRAIN TONNES/HECTARE

***** Tables of means *****

CULTDATE	EARLY	LATER	Mean	
PRIMCULT				
NONE	5.61	5.75	5.68	
DYNDRIVE	5.43	5.83	5.63	
PLOUGH	5.69	5.78	5.73	
TINE	5.56	5.65	5.60	
Mean	5.57	5.75	5.66	
PRSOWCON	GLYPHOS	PARAQUAT	NONE	Mean
PRIMCULT				
NONE	5.74	5.43	5.87	5.68
DYNDRIVE	5.59	5.52	5.77	5.63
PLOUGH	5.73	5.59	5.88	5.73
TINE	5.55	5.48	5.78	5.60
Mean	5.65	5.51	5.83	5.66
PRSOWCON	GLYPHOS	PARAQUAT	NONE	Mean
CULTDATE				
EARLY	5.55	5.40	5.77	5.57
LATER	5.75	5.61	5.89	5.75
Mean	5.65	5.51	5.83	5.66
	PRSOWC	ON GLYPHOS	PARAQUAT	NONE
PRIMCULT	CULTDA			
NONE	EAR	LY 5.72	the second secon	
	LAT	ER 5.76	5.50	5.98
DYNDRIVE	EAR	LY 5.43	5.32	5.55
	LAT	ER 5.76	5.73	5.99
PLOUGH	EAR			5.99
	LAT	ER 5.86	5.71	5.78
TINE	EAR	LY 5.46	5.46	5.75
	LAT	ER 5.64	5.50	5.80

*** Standard errors of differences of means ***

	PRIMCULT	CULTDATE	PRSOWCON	PRIMCULT CULTDATE
	0.083	0.059	0.236	0.118
	PRIMCULT PRSOWCON	CULTDATE	PRIMCULT CULTDATE PRSOWCON	
	0.265	0.246	0.300	
Except when PRIMCULT	comparing means 0.264	with the same	level(s)	of
CULTDATE		0.246		
PRSOWCON	0.140	0.099	0.197	
PRIMCULT.C	ULTDATE		0.298	

90/R/WW/2 FOSTERS CORNER W.WHEAT AFTER S.BARLEY

GRAIN TONNES/HECTARE

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	CV%
BLOCK.WP1	4	0.289	5.1
BLOCK.WP2	14	0.144	2.5
BLOCK.WP1.WP2	28	0.238	4.2

GRAIN MEAN DM% 89.1

PLOT AREA HARVESTED 0.00080