

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1990

[Full Table of Content](#)



90/R/WW/9 Foliar Potassium Nitrate - W. Wheat

Rothamsted Research

Rothamsted Research (1991) *90/R/WW/9 Foliar Potassium Nitrate - W. Wheat* ; Yields Of The Field Experiments 1990, pp 116 - 117 - DOI: <https://doi.org/10.23637/ERADOC-1-42>

90/R/WW/9

WINTER WHEAT

FOLIAR POTASSIUM NITRATE

Object: To study the effects of foliar applications of potassium nitrate and urea on the yield and nutrient composition of w. wheat - Great Harpenden II.

Sponsor: P.B. Barraclough.

Design: 5 randomised blocks of 8 plots.

Whole plot dimensions: 3.0 x 15.0.

Treatments:

FOLIAR N	Foliar nitrogen; all applications were divided equally and applied on two successive days:
NONE	None
K20E	20 kg K as potassium nitrate at GS 41 on 16 and 17 May, 1990
K20EU40E	" " " " " " plus 40 kg N as urea on 16 and 17 May
U40E	40 kg N as urea on 16 and 17 May
K5M	5 kg K as potassium nitrate at GS 55 on 26 and 27 May
K20M	20 kg K " " " " " " 26 and 27 May
K20EK20M	" " " " " " on 16 and 17 May repeated on 26 and 27 May
K20MK20L	" " " " " " on 26 and 27 May repeated GS 71 on 13 and 14 June

NOTE: All plots received N at 131 kg as 'Nitram' applied on 12 Apr, 1990.

Basal applications: Weedkillers: Bromoxynil at 0.19 kg and ioxynil at 0.19 kg with metsulfuron-methyl at 6.0 g in 200 l. Fungicides: Chlorothalonil at 0.50 kg with propiconazole at 0.12 kg in 200 l. Tridemorph at 0.52 kg in 200 l.

Seed: Mercia, sown at 180 kg.

Cultivations, etc.:- Heavy spring-tine cultivated twice: 16 Oct, 1989. Deep-tine cultivated with vibrating tines, disced, rotary harrowed, seed sown: 17 Oct. Weedkillers applied: 24 Apr, 1990. Chlorothalonil with propiconazole applied: 27 May. Tridemorph applied: 26 June. Combine harvested: 7 Aug. Previous crops: W. barley 1988, potatoes 1989.

NOTES: (1) Leaf samples were taken approximately five days after foliar treatment applications to measure N and K contents.
(2) Components of yield were measured.

90/R/WW/9

GRAIN TONNES/HECTARE

***** Tables of means *****

FOLIAR N	
NONE	9.98
K20E	10.29
K20EU40E	10.05
U40E	9.94
K5M	10.26
K20M	10.15
K20EK20M	10.10
K20MK20L	9.92
Mean	10.09

*** Standard errors of differences of means ***

FOLIAR N
0.226

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	28	0.358	3.5
GRAIN MEAN DM%	90.4		

STRAW TONNES/HECTARE

***** Tables of means *****

FOLIAR N	
NONE	6.97
K20E	7.07
K20EU40E	6.60
U40E	6.27
K5M	6.82
K20M	6.89
K20EK20M	6.54
K20MK20L	6.45
Mean	6.70

STRAW MEAN DM% 87.2

PLOT AREA HARVESTED 0.00230