


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED  
RESEARCH

## Yields of the Field Experiments 1990

[Full Table of Content](#)



---

### 90/W/NB/1 Seedbed and Late N - Navy Beans

#### Rothamsted Research

Rothamsted Research (1991) *90/W/NB/1 Seedbed and Late N - Navy Beans* ; Yields Of The Field Experiments 1990, pp 153 - 154 - DOI: <https://doi.org/10.23637/ERADOC-1-42>

90/W/NB/1

NAVY BEANS

SEEDBED AND LATE N

**Object:** To study the effects of seedbed and late nitrogen fertilizer on the growth and yield of navy beans inoculated with Rhizobium - Woburn, Great Hill Bottom I.

**Sponsors:** J.M. Day, M.L. Lovell.

**Design:** 3 randomised blocks of 18 plots.

**Whole plot dimensions:** 1.5 x 6.0.

**Treatments:** All combinations of:-

1. **SD BED N** Nitrogen fertilizer (kg N) as 'Nitro-Chalk' (27% N), applied to seedbed:

20  
40  
60

2. **APP METH** Methods of applying seedbed nitrogen fertilizer:

BCAST Broadcast by hand after drilling  
COM DRLL Combine drilled with seed

3. **FLOWR N** Nitrogen fertilizer (kg N) as 'Nitro-Chalk', applied at flowering on 28 June, 1990:

0  
120

plus four extra treatments :

**EXTRA**

0 0 0 No nitrogen fertilizer (duplicated)  
0 0 120F 120 kg N at flowering (duplicated)  
120S 0 0 120 kg N broadcast to seedbed  
120S 120F 120 kg broadcast to seedbed, repeated at flowering

**Basal applications:** Manures: (0:24:24) at 200 kg. Weedkillers: Paraquat at 0.80 kg ion in 220 l. Trifluralin at 0.84 kg in 220 l. Monolinuron at 0.56 kg in 220 l. Fungicide: Benomyl at 0.55 kg in 220 l. Irrigation: 24 mm (12 mm applied on two occasions).

**Seed:** Albion, sown at 40 seeds per square metre.

**Cultivations, etc.:-** Ploughed: 22 Nov, 1989. Paraquat applied: 3 May, 1990. PK applied: 18 May. Spring-tine cultivated: 21 May. Trifluralin applied and cultivated: 22 May. Seed sown and seedbed N treatments applied: 24 May. Monolinuron applied: 31 May. Irrigation applied: 1 and 20 June. Remaining N treatments applied: 28 June. Fungicide applied 22 Aug. Hand harvested and threshed by stationary combine harvester: 27 Oct. Previous crops: S. wheat 1988, s. barley 1989.

90/W/NB/1

NOTE: Times of flowering were noted:

**GRAIN (AT 85% DRY MATTER) TONNES/HECTARE**

\*\*\*\*\* Tables of means \*\*\*\*\*

APP METH SD BED N	BCAST	COM DRLL	Mean		
20	2.19	2.30	2.25		
40	1.99	1.99	1.99		
60	2.16	2.02	2.09		
Mean	2.11	2.10	2.11		
<b>FLOWR N</b> SD BED N	0	120	Mean		
20	2.17	2.33	2.25		
40	2.19	1.78	1.99		
60	2.21	1.97	2.09		
Mean	2.19	2.03	2.11		
<b>FLOWR N</b> APP METH	0	120	Mean		
BCAST	2.13	2.09	2.11		
COM DRLL	2.25	1.96	2.10		
Mean	2.19	2.03	2.11		
<b>APP METH</b> SD BED N	BCAST	COM DRLL			
	0	120	0		
20	1.97	2.41	2.36		
40	2.17	1.81	2.22		
60	2.26	2.06	2.17		
			1.88		
<b>EXTRA</b>	0 0 0	0 0 120F	120S 0 0	120S 120F	Mean
	2.29	2.11	2.32	2.39	2.25
<b>GRAND MEAN</b>	2.16				

\*\*\* Standard errors of differences of means \*\*\*

SD BED N	APP METH	FLOWR N	SD BED N	APP METH
0.173	0.141	0.141	0.245	
SD BED N	APP METH	SD BED N	EXTRA	
FLOWR N	FLOWR N	APP METH	FLOWR N	
0.245	0.200	0.347	0.347	min.rep
			0.300	max-min
			0.245	max.rep
<b>EXTRA</b>				
min.rep	120S 0 0 v 120S 120F			
max-min	0 0 0 or 0 0 120F v 120S 0 0 or 120S 120F			
max.rep	0 0 0 v 0 0 120F			

GRAIN MEAN DM% \* PLOT AREA HARVESTED 0.00045