

Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1990

[Full Table of Content](#)



90/R/CS/140 Chemical Reference Plots - S. Barley

Rothamsted Research

Rothamsted Research (1991) *90/R/CS/140 Chemical Reference Plots - S. Barley* ; Yields Of The Field Experiments 1990, pp 57 - 60 - DOI: <https://doi.org/10.23637/ERADOC-1-42>

90/R/CS/140

CHEMICAL REFERENCE PLOTS

Object: To study the persistence in soil of agricultural chemicals applied annually, singly and in combination and their effects on soil microflora and on yield of continuous s. barley - Long Hoos V 3.

Sponsors: R.H. Bromilow, A.A. Evans, P.H. Nicholls.

The 17th year, s. barley.

For previous years see 74-89/R/CS/140.

Design: Single replicate of 32 plots.

Whole plot dimensions: 4.06 x 4.57.

Treatments, applied cumulatively every year except as stated:

All combinations of:-

1. **WEEDKLLR** Weedkiller in autumn:
NONE None
GLYPHOS Glyphosate at 1.4 kg to barley stubble each autumn from 1979 to 1984, at 0.72 kg in 1985, at 0.54 kg in 1986, at 1.3 kg in 1987 and at 1.5 kg in 1988 and 1989.
2. **FUNGCIDE[1]** Fungicide in autumn:
NONE None
TRIADIM Triadimefon at 0.25 kg in autumn 1981, 1982, 1984 to 1989, 0.28 kg in autumn 1983
3. **FUNGCIDE[2]** Fungicide in spring:
NONE None
BENOMYL Benomyl at 4 kg to seedbed
4. **INSECTCDE** Insecticide:
NONE None
CHLORFEN Chlorfenvinphos at 2 kg to the seedbed
5. **NEMACIDE** Nematicide:
NONE None
ALDICARB Aldicarb at 6 kg to the seedbed

NOTE: Glyphosate and triadimefon were applied in 220 l on 3 Oct, 1989. Other treatments were applied on 13 Mar, 1990.

Basal applications: Manures: 'Nitram' at 440 kg. Muriate of potash at 520 kg. Weedkillers: Bromoxynil at 0.20 kg, ioxynil at 0.20 kg and mecoprop at 1.6 kg in 220 l.

Seed: Klaxon, seed not dressed, sown at 160 kg.

90/R/CS/140

Cultivations, etc.:- K applied: 24 Aug, 1989. Ploughed: 28 Nov.
 Spring-tine cultivated, N applied: 12 Mar, 1990. Rotary harrowed,
 seed sown: 13 Mar. Rolled: 14 Mar. Weedkillers applied: 23 May.
 Combine harvested: 1 Aug.

GRAIN TONNES/HECTARE

***** Tables of means *****

FUNGCIDE [1]	NONE	TRIADIM	Mean
WEEDKLLR			
NONE	4.45	4.43	4.44
GLYPHOS	4.50	4.41	4.45
Mean	4.47	4.42	4.45
FUNGCIDE [2]	NONE	BENOMYL	Mean
WEEDKLLR			
NONE	4.44	4.44	4.44
GLYPHOS	4.40	4.51	4.45
Mean	4.42	4.48	4.45
FUNGCIDE [2]	NONE	BENOMYL	Mean
FUNGCIDE [1]			
NONE	4.51	4.44	4.47
TRIADIM	4.33	4.52	4.42
Mean	4.42	4.48	4.45
INSTCDE	NONE	CHLORFEN	Mean
WEEDKLLR			
NONE	4.55	4.34	4.44
GLYPHOS	4.46	4.45	4.45
Mean	4.50	4.39	4.45
INSTCDE	NONE	CHLORFEN	Mean
FUNGCIDE [1]			
NONE	4.46	4.49	4.47
TRIADIM	4.54	4.30	4.42
Mean	4.50	4.39	4.45
INSTCDE	NONE	CHLORFEN	Mean
FUNGCIDE [2]			
NONE	4.41	4.43	4.42
BENOMYL	4.60	4.36	4.48
Mean	4.50	4.39	4.45
NEMACIDE	NONE	ALDICARB	Mean
WEEDKLLR			
NONE	4.42	4.47	4.44
GLYPHOS	4.36	4.55	4.45
Mean	4.39	4.51	4.45

90/R/CS/140

GRAIN TONNES/HECTARE

***** Tables of means *****

NEMACIDE	NONE	ALDICARB	Mean
FUNGCIDE [1]			
NONE	4.41	4.54	4.47
TRIADIM	4.36	4.48	4.42
Mean	4.39	4.51	4.45

NEMACIDE	NONE	ALDICARB	Mean
FUNGCIDE [2]			
NONE	4.39	4.44	4.42
BENOMYL	4.38	4.58	4.48
Mean	4.39	4.51	4.45

NEMACIDE	NONE	ALDICARB	Mean
INSCTCDE			
NONE	4.53	4.47	4.50
CHLORFEN	4.24	4.55	4.39
Mean	4.39	4.51	4.45

FUNGCIDE [1]		NONE	TRIADIM		
WEEDKLLR	FUNGCIDE [2]	NONE	BENOMYL	NONE	BENOMYL
NONE		4.42	4.48	4.46	4.40
GLYPHOS		4.60	4.39	4.19	4.63

FUNGCIDE [1]		NONE	TRIADIM		
WEEDKLLR	INSCTCDE	NONE	CHLORFEN	NONE	CHLORFEN
NONE		4.54	4.36	4.56	4.31
GLYPHOS		4.39	4.61	4.53	4.29

FUNGCIDE [2]		NONE	BENOMYL		
WEEDKLLR	INSCTCDE	NONE	CHLORFEN	NONE	CHLORFEN
NONE		4.43	4.45	4.67	4.22
GLYPHOS		4.39	4.40	4.53	4.49

FUNGCIDE [2]		NONE	BENOMYL		
FUNGCIDE [1]	INSCTCDE	NONE	CHLORFEN	NONE	CHLORFEN
NONE		4.44	4.58	4.49	4.39
TRIADIM		4.38	4.27	4.71	4.32

FUNGCIDE [1]		NONE	TRIADIM		
WEEDKLLR	NEMACIDE	NONE	ALDICARB	NONE	ALDICARB
NONE		4.39	4.51	4.44	4.43
GLYPHOS		4.43	4.57	4.28	4.54

FUNGCIDE [2]		NONE	BENOMYL		
WEEDKLLR	NEMACIDE	NONE	ALDICARB	NONE	ALDICARB
NONE		4.54	4.34	4.29	4.59
GLYPHOS		4.25	4.55	4.47	4.56

90/R/CS/140

GRAIN TONNES/HECTARE

***** Tables of means *****

FUNGICIDE [2]		NONE		BENOMYL	
FUNGICIDE [1]	NEMACIDE	NONE	ALDICARB	NONE	ALDICARB
NONE		4.47	4.55	4.35	4.53
TRIADIM		4.31	4.34	4.41	4.62

WEEDKLLR		NONE		CHLORFEN	
	NEMACIDE	NONE	ALDICARB	NONE	ALDICARB
NONE		4.65	4.45	4.19	4.48
GLYPHOS		4.42	4.50	4.29	4.61

FUNGICIDE [1]		NONE		CHLORFEN	
	NEMACIDE	NONE	ALDICARB	NONE	ALDICARB
NONE		4.54	4.39	4.28	4.69
TRIADIM		4.53	4.56	4.20	4.40

FUNGICIDE [2]		NONE		CHLORFEN	
	NEMACIDE	NONE	ALDICARB	NONE	ALDICARB
NONE		4.40	4.42	4.39	4.47
BENOMYL		4.67	4.53	4.09	4.62

*** Standard errors of differences of means ***

Margins of two factor tables	0.061
Two factor tables	0.086
Three factor tables	0.122

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
WP	6	0.173	3.9

GRAIN MEAN DM% 85.4

PLOT AREA HARVESTED 0.00073