


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1991

[Full Table of Content](#)



91/R/CS/361 Control of Stem Nematode - Lucerne

Rothamsted Research

Rothamsted Research (1992) *91/R/CS/361 Control of Stem Nematode - Lucerne* ; Yields Of The Field Experiments 1991, pp 107 - 109 - DOI: <https://doi.org/10.23637/ERADOC-1-46>

91/R/CS/361

CONTROL OF STEM NEMATODE

Object: To study the effects of rates of carbofuran on the control of stem nematode (*Ditylenchus dipsaci*) and on the yield of two varieties of lucerne of different susceptibility - Long Hoos IV 3.

Sponsors: A.G. Whitehead, A.J.F. Nichols.

The first year, lucerne.

Design: 3 randomised blocks of 8 plots.

Whole plot dimensions: 1.2 x 8.8.

Treatments: All combinations of:-

1. **CARBRATE** Rates of carbofuran (kg), applied to seed furrow at sowing: 21 Aug, 1990.

0.00
0.60
1.20
2.30

2. **VARIETY** Varieties and stem nematode susceptibility:

EUROPE Europe - susceptible
EUVER Euver - partially resistant

Basal applications: Weedkillers: 2,4-DB at 1.3 kg in 220 l. Paraquat at 0.40 kg ion in 220 l. Fluazifop-P-butyl at 0.19 kg in 220 l. Irrigation: 5 mm on seven occasions, total 35 mm.

Cultivations, etc.:- Ploughed and rolled: 20 Aug, 1990. Seed sown, cultivated by rotary grubber: 22 Aug. Irrigated: 28, 30 Aug, 3, 10, 13, 20 and 26 Sept. 2,4-DB applied: 14 Dec. Paraquat applied: 7 Mar, 1991. Fluazifop-P-butyl applied: 10 Apr. Cut: 10 July, 13 Aug and 3 Oct. Previous crops: S. peas 1989, w. wheat 1990.

91/R/CS/361

1ST CUT (10/7/91) DRY MATTER TONNES/HECTARE

***** Tables of means *****

CARBRATE	0.00	0.60	1.20	2.30	Mean
VARIETY					
EUROPE	1.19	2.88	2.85	3.54	2.61
EUVER	3.47	3.22	4.36	4.28	3.83
Mean	2.33	3.05	3.61	3.91	3.22

*** Standard errors of differences of means ***

VARIETY	CARBRATE	VARIETY CARBRATE
0.267	0.377	0.533

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	14	0.653	20.3
1ST CUT MEAN DM%	21.0		

2ND CUT (13/8/91) DRY MATTER TONNES/HECTARE

***** Tables of means *****

CARBRATE	0.00	0.60	1.20	2.30	Mean
VARIETY					
EUROPE	1.44	1.88	2.29	3.04	2.16
EUVER	2.87	3.15	3.28	3.50	3.20
Mean	2.16	2.51	2.79	3.27	2.68

*** Standard errors of differences of means ***

VARIETY	CARBRATE	VARIETY CARBRATE
0.178	0.252	0.356

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	14	0.437	16.3
2ND CUT MEAN DM%	17.7		

91/R/CS/361

3RD CUT (3/10/91) DRY MATTER TONNES/HECTARE

***** Tables of means *****

CARBRATE VARIETY	0.00	0.60	1.20	2.30	Mean
EUROPE	1.23	1.40	1.53	1.69	1.46
EUVER	1.99	1.96	2.06	1.86	1.97
Mean	1.61	1.68	1.80	1.77	1.71

*** Standard errors of differences of means ***

VARIETY	CARBRATE	VARIETY CARBRATE
0.064	0.091	0.129

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	14	0.158	9.2

3RD CUT MEAN DM% 20.0

TOTAL OF 3 CUTS DRY MATTER TONNES/HECTARE

***** Tables of means *****

CARBRATE VARIETY	0.00	0.60	1.20	2.30	Mean
EUROPE	3.86	6.15	6.68	8.27	6.24
EUVER	8.33	8.33	9.70	9.64	9.00
Mean	6.09	7.24	8.19	8.95	7.62

*** Standard errors of differences of means ***

VARIETY	CARBRATE	VARIETY CARBRATE
0.472	0.667	0.943

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	14	1.155	15.2

TOTAL OF 3 CUTS MEAN DM% 19.6

PLOT AREA HARVESTED 0.00045