Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readible, or you suspect there are some problems, please let us know and we will correct that.



Yields of the Field Experiments 1991



Full Table of Content

91/R/BW/2 Sowing Dates, Aphids and Bydv - W. Barley

Rothamsted Research

Rothamsted Research (1992) 91/R/BW/2 Sowing Dates, Aphids and Bydv - W. Barley; Yields Of The Field Experiments 1991, pp 126 - 127 - DOI: https://doi.org/10.23637/ERADOC-1-46

91/R/BW/2

WINTER BARLEY

SOWING DATES, APHIDS AND BYDV

Object: To study the relationship of aphid numbers in suction trap samples to crop populations and the incidence of barley yellow dwarf virus (BYDV) on winter barley sown on a range of dates - Highfield IV.

Sponsors: G.M. Tatchell, R.T. Plumb.

Design: 4 randomised blocks of 10 plots.

Whole plot dimensions: 3.0 x 21.0.

Treatments: All combinations of:-

SOWDATE Dates of sowing:

5 SEPT 5 September, 1990
14 SEPT 14 September
24 SEPT 24 September
4 OCT 4 October
17 OCT 17 October

APHICIDE Aphicide:

NONE None

CYPERMET Cypermethrin at 25 g in 300 l on 6 Nov, 1990

NOTES: (1) All SOWDATE treatments were rotary cultivated on 31 July, 1990, ploughed on 8 Aug and rotary harrowed on the day of sowing. SOWDATE 5 SEPT was rotary harrowed again on 5 Sept.

(2) The experiment was netted against birds during June and July.

Basal applications: Manure: 'Nitram' at 460 kg. Weedkillers: Paraquat at 0.40 kg ion with a wetting agent, 'Vassgro' at 0.06 l, in 200 l. Diflufenican at 0.10 kg and isoproturon at 1.0 kg in 300 l. Fungicides: Fenpropimorph at 0.75 kg in 300 l. Prochloraz at 0.40 kg with tridemorph at 0.26 kg in 200 l. Propiconazole at 0.12 kg in 200 l.

Seed: Magie, sown at 140 kg.

Cultivations, etc.:- Paraquat with wetting agent applied: 3 Sept, 1990. Fenpropimorph applied: 14 Nov. Diflufenican and isoproturon applied: 21 Nov. N applied: 26 Mar, 1991. Prochloraz with tridemorph applied: 23 Apr. Propiconazole applied: 23 May. Combine harvested: 5 Aug. Previous crops: W. wheat 1989, w. barley 1990.

NOTES: (1) Aphid numbers were sampled from September to May.

- (2) BYDV was assessed by enzyme-linked immunosorbent assay from November to May and by visual symptoms during May.
- (3) Components of yield were measured.

91/R/BW/2

GRAIN TONNES/HECTARE

***** Tables of means *****

APHICIDE	NONE	CYPERMET	Mean
SOWDATE			
5 SEPT	6.31	7.09	6.70
14 SEPT	7.73	8.09	7.91
24 SEPT	8.17	8.94	8.55
4 OCT	7.74	7.82	7.78
17 OCT	7.44	7.65	7.54
Mean	7.48	7.92	7.70

*** Standard errors of differences of means ***

SOWDATE	APHICIDE	SOWDATE	
		APHICIDE	
0.202	0.128	0.286	

***** Stratum standard errors and coefficients of variation *****

 Stratum
 d.f.
 s.e.
 cv%

 BLOCK.WP
 27
 0.404
 5.2

GRAIN MEAN DM% 85.6

PLOT AREA HARVESTED 0.00204