


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1991

[Full Table of Content](#)



91/R/RAW/2 Varieties and Fungicides - W. Oilseed Rape

Rothamsted Research

Rothamsted Research (1992) *91/R/RAW/2 Varieties and Fungicides - W. Oilseed Rape* ; Yields Of The Field Experiments 1991, pp 142 - 143 - DOI: <https://doi.org/10.23637/ERADOC-1-46>

91/R/RAW/2

WINTER OILSEED RAPE

VARIETIES AND FUNGICIDES

Object: To investigate the effects of fungicides on a range of low glucosinolate varieties - Black Horse I N.

Sponsors: V.J. Church, B.D.L. Fitt.

Design: 4 randomised blocks of 12 plots.

Whole plot dimensions: 3.0 x 21.0.

Treatments: All combinations of:-

1. **VARIETY** Varieties:

CAPRCORN	Capricorn
COBRA	Cobra
ENVOL	Envol
FALCON	Falcon
LIBRAVO	Libravo
SAMOURAI	Samourai

2. **FUNGICIDE** Fungicides:

NONE	None
PR+CA+IP	Prochloraz at 0.45 kg and carbendazim at 0.17 kg in 200 l on 3 Dec, 1990 and 8 Apr, 1991. Iprodione at 0.50 kg in 200 l on 17 June.

Basal applications: Manure: 'Nitram' at 320 kg on two occasions.

Weedkillers: Paraquat at 0.40 kg ion with a wetting agent, 'Vassgro' at 0.06 l, in 200 l. Metazachlor at 0.75 kg in 200 l. Insecticide: Deltamethrin at 6.2 g and on a second occasion at 12.5 g both in 200 l. Desiccant: Diquat at 0.60 kg ion with a wetting agent, 'Vassgro' at 0.52 l, in 520 l.

Seed: Varieties sown at 120 seeds per square metre. Seed dressed with gamma-HCH, thiram and fenpropimorph except for **VARIETY** ENVOL, which was untreated.

Cultivations, etc.:- Heavy spring-tine cultivated: 31 July, 1990.

Paraquat with wetting agent applied, cultivated with rotary grubber: 3 Sept. Rotary harrowed, seed sown: 4 Sept. Metazachlor applied: 2 Oct. First deltamethrin applied: 23 Oct. First N applied: 26 Feb, 1991. Second N applied: 22 Mar. Second deltamethrin applied: 12 Apr. Desiccant with wetting agent applied: 7 Aug. Combine harvested: 13 Aug. Previous crops: W. wheat 1989, w. barley 1990. to post-harvest.

NOTE: Disease assessments were made from November to post-harvest. Plots were assessed for cabbage stem flea beetle damage in October and pollen beetle damage in April. Glucosinolate levels and oil content were measured.

91/R/RAW/2

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

FUNGCIDE VARIETY	NONE	PR+CA+IP	Mean
CAPRCORN	2.13	2.78	2.46
COBRA	1.70	2.52	2.11
ENVOL	1.93	2.68	2.30
FALCON	2.28	2.73	2.50
LIBRAVO	2.64	2.99	2.81
SAMOURAI	1.78	2.67	2.22
Mean	2.08	2.73	2.40

*** Standard errors of differences of means ***

VARIETY	FUNGCIDE	VARIETY FUNGCIDE
0.159	0.092	0.225

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
BLOCK.WP	33	0.319	13.3

GRAIN MEAN DM% 88.5

PLOT AREA HARVESTED 0.00483