


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



ROTHAMSTED
RESEARCH

Yields of the Field Experiments 1991

[Full Table of Content](#)



91/R/RAW/4 N, S and Glucosinolates - W. Oilseed Rape

Rothamsted Research

Rothamsted Research (1992) *91/R/RAW/4 N, S and Glucosinolates - W. Oilseed Rape* ; Yields Of The Field Experiments 1991, pp 144 - 146 - DOI: <https://doi.org/10.23637/ERADOC-1-46>

91/R/RAW/4

WINTER OILSEED RAPE

N, S AND GLUCOSINOLATES

Object: To study the separate and combined effects of rates of nitrogen and sulphur on the quality and yield of three varieties of w. oilseed rape - Pastures.

Sponsors: J.E. Fieldsend, J. Spink, J.E. Leach, H. Stevenson.

Design: 4 replicates of 3 x 3 x 3 in blocks of 9 plots.

Whole plot dimensions: 3.0 x 21.0.

Treatments: All combinations of:

1. **VARIETY** Varieties:

ARIANA	Ariana
FALCON	Falcon
LIBRAVO	Libravo

2. **N** Nitrogen fertilizer (kg N) as 'Nitram' on 12 Mar, 1991:

0
150
250

3. **S** Sulphur (kg S) as calcium sulphate on 13 Mar:

0
50
100

NOTE: Sulphur was applied as gypsum (17.5% S).

Basal applications: Manure: Magnesian limestone at 5.0 t. Weedkiller: Metazachlor at 0.75 kg in 200 l. Fungicide: Prochloraz at 0.40 kg in 300 l. Insecticide: Deltamethrin at 6.2 g in 200 l. Irrigation: 25 mm applied on two occasions.

Seed: Varieties, dressed fenpropimorph, gamma-HCH and thiram, sown at 120 seeds per square metre.

Cultivations, etc.:- Magnesian limestone applied: 31 July, 1990.

Ploughed: 3 Aug. Rotary harrowed: 28 Aug. Rotary harrowed, seed sown: 29 Aug. Weedkiller applied: 30 Aug. Irrigation applied: 13 and 27 Sept. Deltamethrin applied: 15 Oct. Fungicide applied: 23 Apr, 1991. Combine harvested: 8 Aug. Previous crops: W. wheat 1989, w. barley 1990.

NOTE: Crop samples were taken on five occasions throughout the season and a further five during seed development, to measure nitrogen, sulphur and glucosinolate content.

91/R/RAW/4

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Tables of means *****

	N	0	150	250	Mean
VARIETY					
ARIANA		2.94	3.80	3.67	3.47
FALCON		2.87	3.53	3.85	3.42
LIBRAVO		2.88	3.52	3.06	3.15
Mean		2.90	3.61	3.53	3.35
	S	0	50	100	Mean
VARIETY					
ARIANA		3.53	3.39	3.49	3.47
FALCON		3.34	3.48	3.43	3.42
LIBRAVO		3.04	3.27	3.15	3.15
Mean		3.30	3.38	3.35	3.35
	S	0	50	100	Mean
N					
0		2.99	2.82	2.88	2.90
150		3.47	3.69	3.68	3.61
250		3.44	3.64	3.50	3.53
Mean		3.30	3.38	3.35	3.35
	S	0	50	100	
VARIETY	N				
ARIANA	0	3.11	2.81	2.90	
	150	3.73	3.71	3.94	
	250	3.73	3.66	3.61	
FALCON	0	3.00	2.74	2.88	
	150	3.06	3.76	3.76	
	250	3.97	3.94	3.63	
LIBRAVO	0	2.86	2.91	2.85	
	150	3.63	3.58	3.35	
	250	2.63	3.31	3.25	

*** Standard errors of differences of means ***

VARIETY	N	S	VARIETY
	0.067	0.067	0.116
VARIETY	N	VARIETY	N
S	S	N	S
	0.116	0.116	0.211

Except when comparing means with the same level(s) of

VARIETY	0.207
N	0.207
S	0.207
VARIETY . N	0.214
VARIETY . S	0.214
N . S	0.214

91/R/RAW/4

GRAIN (AT 90% DRY MATTER) TONNES/HECTARE

***** Stratum standard errors and coefficients of variation *****

Stratum	d.f.	s.e.	cv%
REP.BLOCK.WP	70	0.283	8.5
GRAIN MEAN DM%	82.8		
SUB PLOT AREA HARVESTED	0.00345		