


Thank you for using eradoc, a platform to publish electronic copies of the Rothamsted Documents. Your requested document has been scanned from original documents. If you find this document is not readable, or you suspect there are some problems, please let us know and we will correct that.



# Yields of the Field Experiments 1991

[Full Table of Content](#)



## 91/R/CS/140 Chemical Reference Plots - S. Barley

### Rothamsted Research

Rothamsted Research (1992) 91/R/CS/140 *Chemical Reference Plots - S. Barley* ; Yields Of The Field Experiments 1991, pp 51 - 54 - DOI: <https://doi.org/10.23637/ERADOC-1-46>

91/R/CS/140

### CHEMICAL REFERENCE PLOTS

**Object:** To study the persistence in soil of agricultural chemicals applied annually, singly and in combination and their effects on soil microflora and on yield of continuous s. barley - Long Hoos V 3.

**Sponsors:** R.H. Bromilow, A.A. Evans, P.H. Nicholls.

The 18th year, s. barley.

For previous years see 74-90/R/CS/140.

**Design:** Single replicate of 32 plots.

**Whole plot dimensions:** 4.06 x 4.57.

Treatments, applied cumulatively every year except as stated:

All combinations of:-

1. **WEEDKLLR** Weedkiller in autumn:

NONE	None
GLYPHOS	Glyphosate at 1.4 kg to barley stubble each autumn from 1979 to 1984, at 0.72 kg in 1985, at 0.54 kg in 1986, at 1.3 kg in 1987 and at 1.5 kg in 1988 to 1990.

2. **FUNGCIDE[1]** Fungicide in autumn:

NONE	None
TRIADIM	Triadimefon at 0.25 kg in autumn 1981, 1982, 1984 to 1990, 0.28 kg in autumn 1983

3. **FUNGCIDE[2]** Fungicide in spring:

NONE	None
BENOMYL	Benomyl at 4 kg to seedbed

4. **INSCTCDE** Insecticide:

NONE	None
CHLORFEN	Chlorfenvinphos at 2 kg to the seedbed

5. **NEMACIDE** Nematicide:

NONE	None
ALDICARB	Aldicarb at 6 kg to the seedbed

**NOTE:** Glyphosate and triadimefon were applied in 220 l on 28 Sept, 1990.  
Other treatments were applied on 8 Apr, 1991.

**Basal applications:** Manures: Muriate of potash at 520 kg. Magnesian limestone at 2.9 t. 'Nitram' at 440 kg. Weedkiller: Fluroxypyr at 0.15 kg in 200 l.

**Seed:** Klaxon, seed not dressed, sown at 160 kg.

91/R/CS/140

**Cultivations, etc.:-** K applied: 14 Sept, 1990. Magnesian limestone applied: 4 Oct. Ploughed: 5 Dec. N applied, heavy spring-tine cultivated, rotary harrowed: 8 Apr, 1991. Seed sown, rolled: 9 Apr. Weedkiller applied: 16 June. Combine harvested: 21 Aug.

**GRAIN TONNES/HECTARE**

\*\*\*\*\* Tables of means \*\*\*\*\*

FUNGCIDE [1]	NONE	TRIADIM	Mean
<b>WEEDKLLR</b>			
NONE	6.07	6.07	6.07
GLYPHOS	6.11	6.09	6.10
Mean	6.09	6.08	6.08
FUNGCIDE [2]	NONE	BENOMYL	Mean
<b>WEEDKLLR</b>			
NONE	6.01	6.13	6.07
GLYPHOS	6.07	6.13	6.10
Mean	6.04	6.13	6.08
FUNGCIDE [2]	NONE	BENOMYL	Mean
FUNGCIDE [1]	NONE	TRIADIM	Mean
NONE	6.04	6.14	6.09
TRIADIM	6.04	6.12	6.08
Mean	6.04	6.13	6.08
INSCTCDE	NONE	CHLORFEN	Mean
<b>WEEDKLLR</b>			
NONE	5.97	6.17	6.07
GLYPHOS	5.95	6.25	6.10
Mean	5.96	6.21	6.08
INSCTCDE	NONE	CHLORFEN	Mean
<b>FUNGCIDE [1]</b>			
NONE	6.02	6.16	6.09
TRIADIM	5.90	6.26	6.08
Mean	5.96	6.21	6.08
INSCTCDE	NONE	CHLORFEN	Mean
<b>FUNGCIDE [2]</b>			
NONE	5.90	6.18	6.04
BENOMYL	6.02	6.24	6.13
Mean	5.96	6.21	6.08
NEMACIDE	NONE	ALDICARB	Mean
<b>WEEDKLLR</b>			
NONE	6.00	6.14	6.07
GLYPHOS	6.14	6.06	6.10
Mean	6.07	6.10	6.08

91/R/CS/140

**GRAIN TONNES/HECTARE**

\*\*\*\*\* Tables of means \*\*\*\*\*

NEMACIDE	NONE	ALDICARB	Mean
<b>FUNGCIDE [1]</b>			
NONE	6.01	6.17	6.09
TRIADIM	6.13	6.03	6.08
Mean	6.07	6.10	6.08
<b>FUNGCIDE [2]</b>			
NONE	6.09	5.99	6.04
BENOMYL	6.05	6.21	6.13
Mean	6.07	6.10	6.08
<b>INSCTCDE</b>			
NONE	5.95	5.96	5.96
CHLORFEN	6.19	6.23	6.21
Mean	6.07	6.10	6.08
<b>WEEDKLLR FUNGCIDE [1]</b>			
	NONE		TRIADIM
	NONE	BENOMYL	NONE
WEEDKLLR	FUNGCIDE [2]		BENOMYL
NONE		5.98	6.05
GLYPHOS		6.10	6.03
<b>WEEDKLLR INSCTCDE</b>			
	NONE		TRIADIM
	NONE	CHLORFEN	NONE
WEEDKLLR	FUNGCIDE [1]		CHLORFEN
NONE		6.05	5.89
GLYPHOS		5.99	6.24
<b>WEEDKLLR INSCTCDE</b>			
	NONE		BENOMYL
	NONE	CHLORFEN	NONE
WEEDKLLR	FUNGCIDE [2]		CHLORFEN
NONE		5.85	6.08
GLYPHOS		5.95	6.19
<b>FUNGCIDE [1] INSCTCDE</b>			
	NONE		BENOMYL
	NONE	CHLORFEN	NONE
FUNGCIDE [1]	INSCTCDE		CHLORFEN
NONE		6.02	6.02
TRIADIM		5.78	6.30
<b>WEEDKLLR NEMACIDE</b>			
	NONE		TRIADIM
	NONE	ALDICARB	NONE
WEEDKLLR	FUNGCIDE [1]		ALDICARB
NONE		5.92	6.08
GLYPHOS		6.10	6.18
<b>WEEDKLLR NEMACIDE</b>			
	NONE		BENOMYL
	NONE	ALDICARB	NONE
WEEDKLLR	FUNGCIDE [2]		ALDICARB
NONE		6.00	6.00
GLYPHOS		6.18	5.95
			6.10
			6.17

91/R/CS/140

GRAIN TONNES/HECTARE

\*\*\*\*\* Tables of means \*\*\*\*\*

FUNGicide [2]		NONE		BENOMYL	
FUNGCIDE [1]	NEMACIDE	NONE	ALDICARB	NONE	ALDICARB
	NONE	6.06	6.02	5.97	6.31
	TRIADIM	6.13	5.95	6.13	6.10
WEEDKLLR		INSCTCDE		CHLORFEN	
WEEDKLLR	NEMACIDE	NONE	ALDICARB	NONE	ALDICARB
	NONE	5.88	6.05	6.12	6.22
	GLYPHOS	6.02	5.88	6.26	6.24
FUNGCIDE [1]		INSCTCDE		CHLORFEN	
FUNGCIDE [1]	NEMACIDE	NONE	ALDICARB	NONE	ALDICARB
	NONE	5.98	6.06	6.05	6.28
	TRIADIM	5.93	5.87	6.33	6.19
FUNGCIDE [2]		INSCTCDE		CHLORFEN	
FUNGCIDE [2]	NEMACIDE	NONE	ALDICARB	NONE	ALDICARB
	NONE	5.93	5.87	6.25	6.11
	BENOMYL	5.97	6.06	6.13	6.36

\*\*\* Standard errors of differences of means \*\*\*

Margins of two factor tables 0.129  
Two factor tables 0.182  
Three factor tables 0.257

\*\*\*\*\* Stratum standard errors and coefficients of variation \*\*\*\*\*

Stratum	d.f.	s.e.	cv%
WP	6	0.364	6.0

GRAIN MEAN DM% 88.8

PLOT AREA HARVESTED 0.00069